

## Recombinant Human MAPK3 293 Cell Lysate

Cat. No. MAPK3-4494HCL Lot. No. (See product label)

### SPECIFICATION

|                            |   |
|----------------------------|---|
| <b>Species</b>             | Human   |
| <b>Source</b>              | HEK293  |
| <b>Description</b>         | Antigen standard for mitogen-activated protein kinase 3 (MAPK3), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.   |
| <b>Components</b>          | This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol). |
| <b>Size</b>                | 0.1 mg  |
| <b>Storage Instruction</b> | Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.   |
| <b>Applications</b>        | ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the   |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

**Gene Name** [MAPK3 mitogen-activated protein kinase 3 \[ Homo sapiens \]](#)

**Official Symbol** MAPK3

**Synonyms** MAPK3; mitogen-activated protein kinase 3; PRKM3; ERK1; p44erk1; p44mapk; MAPK 1; MAP kinase 1; MAP kinase 3; MAP kinase isoform p44; insulin-stimulated MAP2 kinase; mitogen-activated protein kinase 1; extracellular signal-related kinase 1; extracellular signal-regulated kinase 1; microtubule-associated protein 2 kinase; ERT2; ERK-1; P44ERK1; P44MAPK; HS44KDAP; HUMKER1A; p44-ERK1; p44-MAPK; MGC20180;

**Gene ID** [5595](#)

**mRNA Refseq** [NM\\_001040056](#)

**Protein Refseq** [NP\\_001035145](#)

**MIM** [601795](#)

**UniProt ID** [P27361](#)

**Chromosome Location** 16p11.2

**Pathway** ALK1 signaling events, organism-specific biosystem; ARMS-mediated activation, organism-specific biosystem; ATF-2 transcription factor network, organism-specific biosystem; Activated TLR4 signalling, organism-specific biosystem; Activation of the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



AP-1 family of transcription factors, organism-specific biosystem; Acute myeloid leukemia, organism-specific biosystem; Acute myeloid leukemia, conserved biosystem;

**Function**

ATP binding; MAP kinase activity; MAP kinase activity; nucleotide binding; phosphatase binding; phosphotyrosine binding; protein binding; protein serine/threonine kinase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA