

## Recombinant Human MAT2B 293 Cell Lysate

**Cat. No.** MAT2B-4453HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for methionine adenosyltransferase II, beta (MAT2B), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	MAT2B methionine adenosyltransferase II, beta [ Homo sapiens ]
<b>Official Symbol</b>	MAT2B
<b>Synonyms</b>	MAT2B; methionine adenosyltransferase II, beta; methionine adenosyltransferase 2 subunit beta; MATIIbeta; SDR23E1; short chain dehydrogenase/reductase family 23E; member 1; MAT II beta; putative protein product of Nbla02999; dTDP-4-keto-6-deoxy-D-glucose 4-reductase; putative dTDP-4-keto-6-deoxy-D-glucose 4-reductase; short chain dehydrogenase/reductase family 23E, member 1; beta regulatory subunit of methionine adenosyltransferase; TGR; MAT-II; Nbla02999; MGC12237;
<b>Gene ID</b>	27430
<b>mRNA Refseq</b>	NM_182796
<b>Protein Refseq</b>	NP_877725
<b>MIM</b>	605527
<b>UniProt ID</b>	Q9NZL9
<b>Chromosome Location</b>	5q34-q35
<b>Pathway</b>	Biological oxidations, organism-specific biosystem; Cysteine and methionine metabolism, organism-specific biosystem; Cysteine and methionine metabolism, conserved biosystem; Metabolic pathways, organism-specific biosystem; Metabolism,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



organism-specific biosystem; Metabolism of amino acids and derivatives, organism-specific biosystem; Methionine degradation, organism-specific biosystem;

**Function**

dTDP-4-dehydrorhamnose reductase activity; methionine adenosyltransferase regulator activity; nucleotide binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA