

## Recombinant Human MBL2 cell lysate

Cat. No. MBL2-001HCL Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Human MBL2 / MBL / COLEC1 derived in CHO cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
<b>Species</b>	Human
<b>Source</b>	CHO
<b>Preparation method</b>	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
<b>Lysis buffer</b>	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
<b>Quality control Testing</b>	12.5% SDS-PAGE Stained with Coomassie Blue
<b>Recommended Usage</b>	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

**Stability**

Samples are stable for up to twelve months from date of receipt at -80°C

**Storage Buffer**

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

**Storage Instruction**

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

## GENE INFORMATION

**Gene Name**

[MBL2 mannose-binding lectin \(protein C\) 2, soluble \[ Homo sapiens \]](#)

**Official Symbol**

MBL2

**Synonyms**

MBL2; mannose-binding lectin (protein C) 2, soluble; mannose binding lectin (protein C) 2, soluble (opsonic defect) , MBL; mannose-binding protein C; COLEC1; collectin-1; mannan-binding lectin; mannose-binding lectin 2, soluble (opsonic defect); mannose-binding lectin (protein C) 2, soluble (opsonic defect); MBL; MBP; MBP1; MBL2D; MBP-C; HSMBPC; MGC116832; MGC116833;

**Gene ID**

[4153](#)

**mRNA Refseq**

[NM\\_000242](#)

**Protein Refseq**

[NP\\_000233](#)

**MIM**

[154545](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>UniProt ID</b>	P11226
<b>Chromosome Location</b>	10q11.2
<b>Pathway</b>	Complement and coagulation cascades, organism-specific biosystem; Complement and coagulation cascades, conserved biosystem; Complement cascade, organism-specific biosystem; Creation of C4 and C2 activators, organism-specific biosystem; Immune System, organism-specific biosystem; Initial triggering of complement, organism-specific biosystem; Innate Immune System, organism-specific biosystem;
<b>Function</b>	bacterial cell surface binding; calcium-dependent protein binding; eukaryotic cell surface binding; mannose binding; mannose binding; protein binding; receptor binding; sugar binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA