

## Recombinant Human MCM4 293 Cell Lysate

Cat. No. MCM4-4418HCL Lot. No. (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for minichromosome maintenance complex component 4 (MCM4), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	MCM4 minichromosome maintenance complex component 4 [ Homo sapiens ]
<b>Official Symbol</b>	MCM4
<b>Synonyms</b>	MCM4; minichromosome maintenance complex component 4; CDC21, MCM4 minichromosome maintenance deficient 4 ( <i>S. cerevisiae</i> ); DNA replication licensing factor MCM4; CDC54; hCdc21; MGC33310; P1 Cdc21; CDC21 homolog; minichromosome maintenance deficient 4; homolog of <i>S. pombe</i> cell division cycle 21; MCM4 minichromosome maintenance deficient 4; CDC21; P1-CDC21;
<b>Gene ID</b>	4173
<b>mRNA Refseq</b>	NM_005914
<b>Protein Refseq</b>	NP_005905
<b>MIM</b>	602638
<b>UniProt ID</b>	P33991
<b>Chromosome Location</b>	8q12-q13
<b>Pathway</b>	Activation of ATR in response to replication stress, organism-specific biosystem; Activation of the pre-replicative complex, organism-specific biosystem; Assembly of the pre-replicative complex, organism-specific biosystem; C-MYB transcription factor network, organism-specific biosystem; Cell Cycle, organism-specific biosystem; Cell

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Cycle Checkpoints, organism-specific biosystem; Cell Cycle, Mitotic, organism-specific biosystem;

**Function**

ATP binding; contributes\_to ATP-dependent DNA helicase activity; DNA binding; helicase activity; hydrolase activity; nucleotide binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA