

Active Recombinant Human MDM2 protein, His-tagged

Cat. No. MDM2-568H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human MDM2 protein(Q00987)(1-218 aa), fused with N-terminal His tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	1-218 aa
Form	PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	<p>Mouse double minute 2 homolog (MDM2) also known as E3 ubiquitin-protein ligase. Mdm2 is a cellular oncoprotein that recognizes the N-terminal trans-activation domain (TAD) of the p53 tumor suppressor and as an inhibitor of p53 transcriptional activation. The human homologue of this protein is sometimes called Hdm2. The p53 tumor suppressor is the key target of MDM2. It has been identified as a p53 interacting protein that represses p53 transcriptional activity and also acts as an E3 ubiquitin ligase, targeting both itself and p53 for degradation by the proteasome. MDM2 is capable of auto-polyubiquitination, and in complex with p300, a cooperating E3 ubiquitin ligase, is capable of polyubiquitinating p53. Besides, S100 Calcium Binding Protein (S100) has been identified as an interactor of MDM2, thus a binding ELISA assay was conducted to detect the interaction of recombinant human MDM2 and recombinant human S100. Briefly, MDM2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 ul were then transferred to S100-</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-MDM2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 µ stop solution to the wells and read at 450 nm immediately. The binding activity of recombinant human MDM2 and recombinant human S100 was shown, the EC50 is 0.037 ug/ml.

Molecular Mass 28.2kDa

Endotoxin <1.0EU per 1µg (determined by the LAL method).

Purity > 95%

Storage Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.

Reconstitution Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name MDM2 Mdm2, p53 E3 ubiquitin protein ligase homolog (mouse) [Homo sapiens]

Official Symbol MDM2

Synonyms

MDM2; Mdm2, p53 E3 ubiquitin protein ligase homolog (mouse); Mdm2 p53 binding protein homolog (mouse), Mdm2, transformed 3T3 cell double minute 2, p53 binding protein (mouse), mouse double minute 2, human homolog of; p53 binding protein; E3 ubiquitin-protein ligase Mdm2; HDM2; HDMX; MGC5370; oncoprotein Mdm2; MDM2 variant FB28; MDM2 variant FB30; double minute 2, human homolog of; p53-binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

protein; Mdm2, transformed 3T3 cell double minute 2, p53 binding protein; hdm2; ACTFS; MGC71221;

Gene ID 4193


mRNA Refseq NM_002392

Protein Refseq NP_002383


MIM 164785

UniProt ID Q00987

SDS-PAGE



Activity



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA