

## Recombinant Human MEF2C 293 Cell Lysate

Cat. No. MEF2C-4375HCL Lot. No. (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for myocyte enhancer factor 2C (MEF2C), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	MEF2C myocyte enhancer factor 2C [ Homo sapiens ]
<b>Official Symbol</b>	MEF2C
<b>Synonyms</b>	MEF2C; myocyte enhancer factor 2C; myocyte-specific enhancer factor 2C; MADS box transcription enhancer factor 2, polypeptide C; C5DELq14.3; DEL5q14.3;
<b>Gene ID</b>	4208
<b>mRNA Refseq</b>	NM_001131005
<b>Protein Refseq</b>	NP_001124477
<b>MIM</b>	600662
<b>UniProt ID</b>	Q06413
<b>Chromosome Location</b>	5q14
<b>Pathway</b>	Activated TLR4 signalling, organism-specific biosystem; Adipogenesis, organism-specific biosystem; CDO in myogenesis, organism-specific biosystem; Circadian Clock, organism-specific biosystem; Delta-Notch Signaling Pathway, organism-specific biosystem; Developmental Biology, organism-specific biosystem; ERK/MAPK targets, organism-specific biosystem;
<b>Function</b>	AT DNA binding; DNA binding; DNA binding; RNA polymerase II core promoter

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



proximal region sequence-specific DNA binding transcription factor activity involved in positive regulation of transcription; RNA polymerase II core promoter sequence-specific DNA binding transcription factor activity; RNA polymerase II regulatory region sequence-specific DNA binding; activating transcription factor binding; miRNA binding; protein binding; protein heterodimerization activity; sequence-specific DNA binding RNA polymerase II transcription factor activity; sequence-specific DNA binding transcription factor activity; transcription regulatory region DNA binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA