

## Active Recombinant Human MET Mutant (Y1230C), GST-tagged

Cat. No. MET-69H Lot. No. (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant human MET (Y1230C) (956-end) was expressed by baculovirus in Sf9 insect cells using an N-terminal GST tag.
<b>Species</b>	Human
<b>Source</b>	Sf9 Cells
<b>ProteinLength</b>	956-end a.a.
<b>Description</b>	<p>MET is a proto-oncogene that encodes a transmembrane growth factor receptor which is a heterodimer of two disulphide linked chains of 50 kDa (alpha) and 145 kDa (beta). MET is widely expressed in the kidney, brain, lung, skin, and embryonic tissue. Hepatocyte growth factor (HGF) binds to MET and activates its tyrosine kinase activity. MET is overexpressed and activated in a variety of human cancers including pancreatic, colon, gastric, cervical and ovarian cancers and has been shown to be involved in tumor cell migration and invasion. MET(P991S) is one of the native mutant forms of MET.</p>
<b>Form</b>	Recombinant protein stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol.
<b>Bio-activity</b>	The specific activity was determined to be 260 nmol/min/mg
<b>Molecular Mass</b>	~81 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Purity</b>	>85% by densitometry.
<b>Applications</b>	Kinase Assay
<b>Storage</b>	Store product at –70oC. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.
<b>Concentration</b>	0.1 µg/µl

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">MET met proto-oncogene (hepatocyte growth factor receptor) [ Homo sapiens ]</a>
<b>Official Symbol</b>	MET
<b>Synonyms</b>	MET; met proto-oncogene (hepatocyte growth factor receptor); hepatocyte growth factor receptor; HGFR; RCCP2; SF receptor; HGF receptor; oncogene MET; HGF/SF receptor; proto-oncogene c-Met; scatter factor receptor; tyrosine-protein kinase Met; met proto-oncogene tyrosine kinase; AUTS9; c-Met;
<b>Gene ID</b>	<a href="#">4233</a>
<b>mRNA Refseq</b>	<a href="#">NM_000245</a>
<b>Protein Refseq</b>	<a href="#">NP_000236</a>
<b>MIM</b>	<a href="#">164860</a>
<b>UniProt ID</b>	<a href="#">P08581</a>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Chromosome Location</b>	7q31
<b>Pathway</b>	Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Alpha6-Beta4 Integrin Signaling Pathway, organism-specific biosystem; Arf6 signaling events, organism-specific biosystem; Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem; Axon guidance, organism-specific biosystem;
<b>Function</b>	ATP binding; hepatocyte growth factor-activated receptor activity; nucleotide binding; protein binding; protein tyrosine kinase activity; protein tyrosine kinase activity; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA