

Active Recombinant Human MMP3 protein, His-tagged

Cat. No. MMP3-1123H Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human MMP3 protein(P08254)(18~477 aa), fused with N-terminal His tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	18-477 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	<p>Matrix metalloproteinases are a family of zinc and calcium dependent endopeptidases with the combined ability to degrade all the components of the extracellular matrix. MMP-3 (stromelysin-1), can degrade a broad range of substrates including collagen alpha chains, aggrecan, laminin, fibronectin, and elastin. MMP-3 does not cleave the triple helical region of interstitial collagens, a characteristic which distinguishes the stromelysins from the collagenases. MMP-3 is expressed by fibroblasts, chondrocytes, osteoblasts, endothelial cells, smooth muscle cells and macrophages. Structurally, MMP-3 may be divided into several distinct domains; a pro-domain which is cleaved upon activation; a catalytic domain containing the zinc binding site; a short hinge region and a carboxyl terminal (hemopexin-like) domain. The activity of recombinant human MMP-3 is measured by its ability to cleave a fluorogenic peptide substrate Mca-Arg-Pro-Lys-Pro-Val-Glu-Nval-Trp-Arg-Lys(Dnp)-NH₂ in the assay buffer 50 mM Tris, 10 mM Ca</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Cl₂, 150 mM NaCl, 0.05% (w/v) Brij-35, pH 7.5. The rhMMP3 is diluted to 200 ug/ml in assay buffer, then activated with 5 ug/ml chymotrypsin at 37°C for 30min followed by adding 2 mM PMSF to stop activation. The activated rhMMP3 is diluted to 6.25 ug/mL in assay buffer. Loading into a black well plate 50 μ of 6.25 ug/mL rhMMP3 and start the reaction by adding 50 μ of 20 μM substrate, with a substrate blank containing 50 μ assay buffer, 50 μ substrate, and no rhMMP3. Then read at excitation and emission wavelengths of 320 nm and 405 nm, respectively, in kinetic mode for 5 minutes. The specific activity of recombinant human MMP3 is > 90 pmol/min/g.

Molecular Mass	53.8kDa
Endotoxin	<1.0EU per 1μg (determined by the LAL method).
Purity	> 95% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION


Gene Name	MMP3 matrix metalloproteinase 3 (stromelysin 1, progelatinase) [Homo sapiens]
Official Symbol	MMP3
Synonyms	MMP3; matrix metalloproteinase 3 (stromelysin 1, progelatinase); matrix metalloproteinase 3 (stromelysin 1, progelatinase), STMY, STMY1; stromelysin-1; transin-1; proteoglycanase; matrix metalloproteinase-3; matrix metalloproteinase 3 (stromelysin 1, progelatinase); SL-1; STMY; STR1; CHDS6; MMP-3; STMY1;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MGC126102; MGC126103; MGC126104;

Gene ID 4314**mRNA Refseq** NM_002422**Protein Refseq** NP_002413**MIM** 185250**UniProt ID** P08254**SDS-PAGE** Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA