

## Recombinant Human MS4A2 293 Cell Lysate

**Cat. No.** MS4A2-4126HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for membrane-spanning 4-domains, subfamily A, member 2 (Fc fragment of IgE, high affinity I, receptor for; beta polypeptide) (MS4A2), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

**Gene Name** [MS4A2 membrane-spanning 4-domains, subfamily A, member 2 \[ Homo sapiens \]](#)

**Official Symbol** [MS4A2](#)

**Synonyms**

MS4A2; membrane-spanning 4-domains, subfamily A, member 2; APY, FCER1B, IgE responsiveness (atopic) , IGER, membrane spanning 4 domains, subfamily A, member 2 (Fc fragment of IgE, high affinity I, receptor for; beta polypeptide); high affinity immunoglobulin epsilon receptor subunit beta; Fc fragment of IgE; high affinity I; receptor for; beta polypeptide; MS4A1; igE Fc receptor subunit beta; high affinity IgE receptor beta subunit; immunoglobulin E receptor, high affinity, beta polypeptide; Fc fragment of IgE, high affinity I, receptor for; beta polypeptide; High affinity immunoglobulin epsilon receptor beta-subunit (FcERI) (IgE Fc receptor, beta-subunit) (Fc epsilon receptor I beta-chain); APY; IGEL; IGER; ATOPY; FCERI; IGHF; FCER1B;

**Gene ID** [2206](#)

**mRNA Refseq** [NM\\_000139](#)

**Protein Refseq** [NP\\_000130](#)

**MIM** [147138](#)

**UniProt ID** [Q01362](#)

**Chromosome** [11q12-q13](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Location</b>	
<b>Pathway</b>	Asthma, organism-specific biosystem; Asthma, conserved biosystem; Fc epsilon RI signaling pathway, organism-specific biosystem; Fc epsilon RI signaling pathway, conserved biosystem; Fc-epsilon receptor I signaling in mast cells, organism-specific biosystem;
<b>Function</b>	IgE binding; SH2 domain binding; calcium channel activity; phosphoprotein binding; protein kinase binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA