

Active Recombinant Full Length Human MYC Protein, C-Flag-tagged

Cat. No. MYC-757HFL Lot. No. (See product label)

SPECIFICATION

Product Overview

Recombinant Full Length Human MYC Protein, fused to Flag-tag at C-terminus, was expressed in Mammalian cells.

Species

Human

Source

Mammalian Cells

Description

This gene is a proto-oncogene and encodes a nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. The encoded protein forms a heterodimer with the related transcription factor MAX. This complex binds to the E box DNA consensus sequence and regulates the transcription of specific target genes. Amplification of this gene is frequently observed in numerous human cancers. Translocations involving this gene are associated with Burkitt lymphoma and multiple myeloma in human patients. There is evidence to show that translation initiates both from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site, resulting in the production of two isoforms with distinct N-termini.

Form

25 mM Tris HCl, pH 7.3, 100 mM glycine, 10% glycerol.

Bio-activity

ELISA binding assay
EMSA assay
ELISA capture for autoantibodies

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	50.4 kDa
AA Sequence	<p>LDFFRVVENQQPPATMPLNVSFTNRNYDLDYDSVQPYFYCDEEENFYQQQQQSEL QPPAPSEDIWKKFEL LPTPPLSPSRRSGLCSPSYVAVTPFSLRGDNDGGGGSFSTA DQLEMVTELLGGDMVNSFCIDPDETFI KNIIQDCMWSGFSAAAKLVSEKLASYQ AARKDSGSPNPARGHSVCSTSSLYLQDLSAAAASECIDPSVVF PYPLNDSSSPKSCA SQDSSAFSPSSDILLSSTESSPQGSPEPLVLHEETPPTTSSDSEEEQEDEEEEIDVV S VEKRQAPGKRSESGSPSAGGHKPPHSPLVLKRCHVSTHQHNYAAPPSTRKDYP AKRVKLDVSVLRQ ISNNRKCTSPRSSDTEENVKRRTHNVLERQRRNELKRSFFAL RDQIPELENNEKAPKVVILKATAYILS VQAEQKLISEEDLLRKRREQLKHKLEQLRNSCATRTRPLEQKLISEEDLAANDILDY KDDDDKV</p>
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining.
Stability	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
Storage	Store at -80 centigrade.
Concentration	>50 ug/mL as determined by microplate BCA method.
Preparation	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Protein Families	Druggable Genome, Embryonic stem cells, Induced pluripotent stem cells, Stem cell - Pluripotency, Stem cell relevant signaling - JAK/STAT signaling pathway, Stem cell relevant signaling - TGFb/BMP signaling pathway, Stem cell relevant signaling - Wnt Signaling pathway, Transcription Factors

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Pathways Acute myeloid leukemia, Bladder cancer, Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, ErbB signaling pathway, Jak-STAT signaling pathway, MAPK signaling pathway, Pathways in cancer, Small cell lung cancer, TGF-beta signaling pathway, Thyroid cancer, Wnt signaling pathway

Full Length Full L.

GENE INFORMATION

Gene Name MYC MYC proto-oncogene, bHLH transcription factor [Homo sapiens (human)]

Official Symbol MYC

Synonyms MRTL; MYCC; c-Myc; bHLHe39

Gene ID 4609

mRNA Refseq NM_002467.6

Protein Refseq NP_002458.2


MIM 190080

UniProt ID P01106

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Coomassie blue staining of purified MYC protein.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA