

Myc CoIP Agarose kit

Cat. No. Myc-03A Lot. No. (See product label)

SPECIFICATION


Product Overview	<p>The Myc-Trap Agarose Kit contains Myc-Trap Agarose, lysis, wash, and elution buffers for efficient immunoprecipitation of Myc-tagged proteins and their interacting factors.</p> <ul style="list-style-type: none"> • Fast, reliable & efficient one-step immunoprecipitation • Ready-to-use • No heavy & light antibody chains on the SDS-PAGE • Stable under harsh washing conditions • Elution with peptide
Specificity/Target	Myc-tag sequence motif EQKLISEEDL at the N-terminus, C-terminus, or internal site of the fusion protein. Endogenous c-myc is NOT bound.
Binding capacity	17.5 µg of recombinant Myc-tagged protein (~42 kDa) per 25 µL bead slurry
Conjugate	Agarose beads; bead size: ~ 90 µm (cross-linked 4 % agarose beads)
Elution buffer	<p>2x Myc peptide.</p> <p>SDS sample buffer</p> <p>0.2 M glycine pH 2.5</p>
Wash buffer compatibility	2 M NaCl, 10 mM DTT, 2 % Nonidet P40 Substitute, 1 % Triton X-100
Type	Nanobody

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Affinity (KD)	1x Myc-tag:Dissociation constant KD of 500 nM 2x Myc-tag:Dissociation constant KD of 0.5 Nm
Storage Buffer	20% ethanol
Applications	IP, CoIP, ChIP, RIP



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA