

Recombinant Human NCF1 293 Cell Lysate

Cat. No. NCF1-3950HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for neutrophil cytosolic factor 1 (NCF1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [NCF1 neutrophil cytosolic factor 1 \[Homo sapiens \]](#)

Official Symbol NCF1

Synonyms

NCF1; neutrophil cytosolic factor 1; neutrophil cytosolic factor 1 (47kD, chronic granulomatous disease, autosomal 1); neutrophil cytosol factor 1; chronic granulomatous disease; autosomal 1; NADPH oxidase organizer 2; NCF1A; NOXO2; p47phox; SH3PXD1A; NCF-1; NCF-47K; p47-phox; nox organizer 2; nox-organizing protein 2; 47 kDa neutrophil oxidase factor; neutrophil NADPH oxidase factor 1; SH3 and PX domain-containing protein 1A; 47 kDa autosomal chronic granulomatous disease protein; neutrophil cytosolic factor 1, (chronic granulomatous disease, autosomal 1); FLJ79451;

Gene ID [653361](#)

mRNA Refseq [NM_000265](#)

Protein Refseq [NP_000256](#)

MIM [608512](#)

UniProt ID [P14598](#)

Chromosome Location 7q11.23

Pathway Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



pathway, conserved biosystem; Fc gamma R-mediated phagocytosis, organism-specific biosystem; Fc gamma R-mediated phagocytosis, conserved biosystem; Leishmaniasis, organism-specific biosystem; Leishmaniasis, conserved biosystem; Leukocyte transendothelial migration, organism-specific biosystem;

Function

GTP binding; GTPase activity; SH3 domain binding; electron carrier activity; phosphatidylinositol binding; protein binding; superoxide-generating NADPH oxidase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA