

Recombinant Human NCK1 293 Cell Lysate

Cat. No. NCK1-3947HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for NCK adaptor protein 1 (NCK1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	NCK1 NCK adaptor protein 1 [Homo sapiens]
Official Symbol	NCK1
Synonyms	NCK1; NCK adaptor protein 1; NCK; cytoplasmic protein NCK1; NCKalpha; NCK tyrosine kinase; melanoma NCK protein; SH2/SH3 adaptor protein NCK-alpha; non-catalytic region of tyrosine kinase; nck-1; MGC12668;
Gene ID	4690
mRNA Refseq	NM_001190796
Protein Refseq	NP_001177725
MIM	600508
UniProt ID	P16333
Chromosome Location	3q21
Pathway	Activation of Rac, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem; Angiotensin receptor Tie2-mediated signaling, organism-specific biosystem; Antigen Activates B Cell Receptor Leading to Generation of Second Messengers, organism-specific biosystem; Arf6 signaling events, organism-specific biosystem; Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

cytoskeletal adaptor activity; protein binding; protein domain specific binding; receptor binding; receptor signaling complex scaffold activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA