

Recombinant Human NCOA3 293 Cell Lysate

Cat. No. NCOA3-3941HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for nuclear receptor coactivator 3 (NCOA3), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	NCOA3 nuclear receptor coactivator 3 [Homo sapiens]
Official Symbol	NCOA3
Synonyms	NCOA3; nuclear receptor coactivator 3; ACTR; AIB1; bHLHe42; CAGH16; KAT13B; p/CIP; RAC3; receptor associated coactivator 3; SRC 3; SRC3; thyroid hormone receptor activator molecule 1; TNRC16; TRAM 1; CBP-interacting protein; receptor-associated coactivator 3; amplified in breast cancer 1 protein; steroid receptor coactivator protein 3; class E basic helix-loop-helix protein 42; pCIP; AIB-1; CTG26; SRC-3; TNRC14; TRAM-1; MGC141848;
Gene ID	8202
mRNA Refseq	NM_001174087
Protein Refseq	NP_001167558
MIM	601937
UniProt ID	Q9Y6Q9
Chromosome Location	20q12
Pathway	Androgen Receptor Signaling Pathway, organism-specific biosystem; BMAL1:CLOCK/NPAS2 Activates Circadian Expression, organism-specific biosystem; Circadian Clock, organism-specific biosystem; Circadian Repression of Expression by

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



REV-ERBA, organism-specific biosystem; Developmental Biology, organism-specific biosystem; FOXA1 transcription factor network, organism-specific biosystem; Fatty acid, triacylglycerol, and ketone body metabolism, organism-specific biosystem;

Function

androgen receptor binding; chromatin binding; estrogen receptor binding; histone acetyltransferase activity; ligand-dependent nuclear receptor binding; ligand-dependent nuclear receptor transcription coactivator activity; nuclear hormone receptor binding; protein N-terminus binding; protein binding; signal transducer activity; thyroid hormone receptor binding; transcription coactivator activity; transcription coactivator activity; transferase activity, transferring acyl groups;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA