

Recombinant Human NEDD4 293 Cell Lysate

Cat. No. NEDD4-3886HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for neural precursor cell expressed, developmentally down-regulated 4 (NEDD4), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	NEDD4 neural precursor cell expressed, developmentally down-regulated 4, E3 ubiquitin protein ligase [Homo sapiens]
Official Symbol	NEDD4
Synonyms	NEDD4; neural precursor cell expressed, developmentally down-regulated 4, E3 ubiquitin protein ligase; neural precursor cell expressed, developmentally down regulated 4; E3 ubiquitin-protein ligase NEDD4; KIAA0093; MGC176705; NEDD4 1; receptor potentiating factor 1; RPF1; NEDD-4; receptor-potentiating factor 1; cell proliferation-inducing gene 53 protein; neural precursor cell expressed developmentally down-regulated protein 4; NEDD4-1;
Gene ID	4734
mRNA Refseq	NM_006154
Protein Refseq	NP_006145
MIM	602278
UniProt ID	P46934
Chromosome Location	15q
Pathway	Adaptive Immune System, organism-specific biosystem; Antigen processing: Ubiquitination and Proteasome degradation, organism-specific biosystem; Antiviral

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



mechanism by IFN-stimulated genes, organism-specific biosystem; Class I MHC mediated antigen processing & presentation, organism-specific biosystem; Cytokine Signaling in Immune system, organism-specific biosystem;

Function

RNA polymerase binding; acid-amino acid ligase activity; beta-2 adrenergic receptor binding; ligase activity; phosphoserine binding; phosphothreonine binding; proline-rich region binding; proline-rich region binding; protein binding; protein domain specific binding; sodium channel inhibitor activity; ubiquitin binding; ubiquitin-protein ligase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA