

Recombinant Human NFATC1 Protein, Full Length, C-Flag tagged

Cat. No. NFATC1-20HFL Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant protein of human nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1 (NFATC1), transcript variant 1 with a C-Flag tag was expressed in HEK293T.

Species Human

Source HEK293

ProteinLength Full Length

Description The product of this gene is a component of the nuclear factor of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation, and an inducible nuclear component. Proteins belonging to this family of transcription factors play a central role in inducible gene transcription during immune response. The product of this gene is an inducible nuclear component. It functions as a major molecular target for the immunosuppressive drugs such as cyclosporin A. Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. Different isoforms of this protein may regulate inducible expression of different cytokine genes.

Molecular Mass 77.6 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence

MPSTSFVPVPSKFPLGPAAAVFGRGETLGPAPRAGGTMKSAEEEHYGYASSNVSPAL
 PLPTAHSTLPAPCHNLQSTSTPGIIPADHPSGYGAALDGGPAGYFLSSGHTRPDGAP
 ALESPRIEITSCGLYHNNNQFFHDVEVEDVLPSSKRSPSTATLSLPSLEAYRDPSCL
 SPASSLSSRSCNSEASSYESNYSYPYASPQTSPWQSPCVSPKTTDPEEGFPRGLG
 ACTLLGSPRHSPSTSPRASVTEESWLGARSSRPASPCNKRKYSLNGRQPPYSPHH
 SPTSPHGS PRVSVTDDSWLGNTTQYTSSAIVAINALTTDSSLDLGDGVPVKSRT
 TLEQPPSVALKVEPVGEDLGSPPPPADFAPEDYSSFQHIRKGGFCDQYLAVPQHYPY
 QWAKPKPLSPTSYSMSPTLPALDWQLPSHSGPYELRIEVQPKSHHRAHYETEGSRG
 AVKASAGGHPIVQLHGYLENEPLMLQLFIGTADDRLLRPHAFYQVHRITGKTVSTTSH
 EAILSNTKVL EIPLLPENSMRAVIDCAGILKLRNSDIELRKGETDIGRKNTRVRLVFRVH
 VPQPSGRTL SLQVASNPIECSQRSAQELPLVEKQSTDSYPVVGKMMVLSGHNFLQ
 DSKVIFVEKAPDGHHVWEMEAKTDRDLCKPNSLVVEIPFRNQRITSPVHVSFYVCN
 GKRKRSQYQRFTYLPANGNAIFLTVSREHERVGCFFSGPTRTRPLEQKLISEEDLAA
 NDILDYKDDDDKV

Purity

> 80% as determined by SDS-PAGE and Coomassie blue staining

Notes

For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.

Stability

Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.

Storage

Store at -80 centigrade.

Concentration

> 0.05 µg/µL as determined by microplate BCA method

Storage Buffer

25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol


GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	NFATC1 nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1 [Homo sapiens (human)]
Official Symbol	NFATC1
Synonyms	NFATC1; nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1; nuclear factor of activated T-cells, cytoplasmic 1; NF ATC; NFAT2; NFATc; NF-ATc1; NFAT transcription complex cytosolic component; NF-ATC; MGC138448;
Gene ID	4772
mRNA Refseq	NM_172390
Protein Refseq	NP_765978
MIM	600489
UniProt ID	O95644

SDS-PAGE


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA