

Recombinant Human NFATC2 protein, MYC/DDK-tagged

Cat. No. NFATC2-182H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human NFATC2, transcript variant 2, fused with MYC/DDK tag at C-terminal was expressed in HEK293.
Species	Human
Source	HEK293
Description	This gene is a member of the nuclear factor of activated T cells (NFAT) family. The product of this gene is a DNA-binding protein with a REL-homology region (RHR) and an NFAT-homology region (NHR). This protein is present in the cytosol and only translocates to the nucleus upon T cell receptor (TCR) stimulation, where it becomes a member of the nuclear factors of activated T cells transcription complex. This complex plays a central role in inducing gene transcription during the immune response. Alternate transcriptional splice variants encoding different isoforms have been characterized.
Form	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.
Molecular Mass	100 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration	>50 ug/mL as determined by microplate BCA method

GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	NFATC2 nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2 [Homo sapiens]
Official Symbol	NFATC2
Synonyms	NFATC2; nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2; nuclear factor of activated T-cells, cytoplasmic 2; NF ATP; NFAT1; NFATp; NF-ATc2; NFAT pre-existing subunit; T cell transcription factor NFAT1; T-cell transcription factor NFAT1; NFAT transcription complex, preexisting component; preexisting nuclear factor of activated T-cells 2; nuclear factor of activated T-cells, preexisting component; NFATP;
Gene ID	4773
mRNA Refseq	NM_173091
Protein Refseq	NP_775114
MIM	600490
UniProt ID	Q13469
Chromosome Location	20q13.2
Pathway	Axon guidance, organism-specific biosystem; Axon guidance, conserved biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem; B cell receptor signaling pathway, organism-specific biosystem; B cell receptor signaling pathway, conserved biosystem; Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem; Calcium signaling in the CD4+ TCR pathway, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

DNA binding; protein binding; sequence-specific DNA binding; sequence-specific DNA binding transcription factor activity; transcription regulatory region DNA binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA