

Recombinant Human NFATC2 Protein, Myc/DDK-tagged, C13 and N15-labeled

Cat. No. NFATC2-3619H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

NFATC2 MS Standard C13 and N15-labeled recombinant protein (NP_775114) with a C-terminal MYC/DDK tag, was expressed in HEK293 cells.

Species

Human

Source

HEK293

Description

This gene is a member of the nuclear factor of activated T cells (NFAT) family. The product of this gene is a DNA-binding protein with a REL-homology region (RHR) and an NFAT-homology region (NHR). This protein is present in the cytosol and only translocates to the nucleus upon T cell receptor (TCR) stimulation, where it becomes a member of the nuclear factors of activated T cells transcription complex. This complex plays a central role in inducing gene transcription during the immune response. Alternate transcriptional splice variants encoding different isoforms have been characterized.

Molecular Mass

100 kDa

AA Sequence

MNAPERQPPDGGDAPGHEPGGSPQDELDFSILFDYEYLNPNEEEEPNAHKVASPP
SGPAYPDDVLDYGLKPYSPLASLSGEPGRFGEPDRVGPQKFLSAAKPAGASGLSP
RIEITPSHELIVAVGPLRMRDAGLLVEQPPLAGVAASPRFTLPVPGFEGYREPLCLSP
ASSGSSASFISDTFSPYTSPCVSPNNGGPDLLCPQFQNIPIAHYSPRTSPIMSPRTSL
AEDSCLGRHSPVPRPASRSSSPGAKRRHS CAEALVALPPGASPQRSRSRSPSPQPSS

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

HVAPQDHGSPAGYPPVAGSAVIMDALNSLATDSPCGIPPKMWKTSPDPSPVSAAPS
 KAGLPRHIYPAVEFLGPCEQGERRNSAPESILLVPPTWPKPLVPAIPICSIPVTASLPP
 LEWPLSSQSGSYELRIEVQPKPHHRAHYETEGSRGAVKAPTGGHPVVQLHGYMEN
 KPLGLQIFIGTADERILKPHAFYQVHRITGKTVTTTSYEKIVGNTKVLEIPLEPKNNMRA
 TIDCAGILKLRNADIELRKGETDIGRKNTRVRLVFRVHIPESSGRIVSLQTASNPIECSQ
 RSAHELPMVERQDTSCLVYGGQQMILTGQNFTSESKVVFEKTTDGGQIWEA
 TVDKDKSQPNMLFVEIPEYRNKHIRTVPKVNIFYVINGKRKRSQPQHFTYHPVPAIKTE
 PTDEYDPTLICSPTHGGLGSQPYYPQHMPVAESPCLVATMAPCQQFRTGLSSPDA
 RYQQQNPAAVLYQRSKSLSPSLGYYQPALMAAPLSLADAHRSVLVHAGSQGQSS
 ALLHPSPTNQQASPVIIHYSPTNQQLRCSHQEFQHIMYCENFAPGTTTRPGPPPVSQ
 GQRLSPGSYPTVIQQNATSQRAAKNGPPVSDQKEVLPAGVTIKQEQLDQTYLDD
 VNEIIRKEFSGPPARNQTTTRTRPLEQKLISEEDLAANDILDYKDDDDKV

Purity > 80% as determined by SDS-PAGE and Coomassie blue staining

Stability Stable for 3 months from receipt of products under proper storage and handling conditions.

Storage Store at -80 centigrade. Avoid repeated freeze-thaw cycles.

Concentration 50 µg/mL as determined by BCA

Storage Buffer 100 mM glycine, 25 mM Tris-HCl, pH 7.3.

GENE INFORMATION

Gene Name NFATC2 nuclear factor of activated T cells 2 [Homo sapiens (human)]

Official Symbol NFATC2

Synonyms NFATC2; nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

nuclear factor of activated T-cells, cytoplasmic 2; NF ATP; NFAT1; NFATp; NF-ATc2; NFAT pre-existing subunit; T cell transcription factor NFAT1; T-cell transcription factor NFAT1; NFAT transcription complex, preexisting component; preexisting nuclear factor of activated T-cells 2; nuclear factor of activated T-cells, preexisting component; NFATP;

Gene ID [4773](#)


mRNA Refseq [NM_173091](#)

Protein Refseq [NP_775114](#)

MIM [600490](#)

UniProt ID [Q13469](#)

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA