

## Recombinant Human NOXA1 293 Cell Lysate

Cat. No. NOXA1-3749HCL Lot. No. (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for NADPH oxidase activator 1 (NOXA1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	<a href="#">NOXA1 NADPH oxidase activator 1 [ Homo sapiens ]</a>
<b>Official Symbol</b>	NOXA1
<b>Synonyms</b>	NOXA1; NADPH oxidase activator 1; SDCCAG31, serologically defined colon cancer antigen 31; FLJ25475; NY CO 31; p51-nox; Nox activator 1; antigen NY-CO-31; NCF2-like protein; p67phox-like factor; inhibitory NADPH oxidase activator 1; serologically defined colon cancer antigen 31; p51NOX; NY-CO-31; SDCCAG31; MGC131800;
<b>Gene ID</b>	<a href="#">10811</a>
<b>mRNA Refseq</b>	<a href="#">NM_001256067</a>
<b>Protein Refseq</b>	<a href="#">NP_001242996</a>
<b>MIM</b>	<a href="#">611255</a>
<b>UniProt ID</b>	<a href="#">Q86UR1</a>
<b>Chromosome Location</b>	9q34.3
<b>Pathway</b>	RAC1 signaling pathway, organism-specific biosystem;
<b>Function</b>	Rac GTPase binding; SH3 domain binding; enzyme binding; protein binding; superoxide-generating NADPH oxidase activator activity; superoxide-generating

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



NADPH oxidase activator activity;

☎ Tel: 1-631-559-9269 1-516-512-3133

☎ Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) ☎ Fax: 1-631-938-8127

☎ 45-1 Ramsey Road, Shirley, NY 11967, USA