

## Recombinant Human NR0B1 293 Cell Lysate

**Cat. No.** NR0B1-3723HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for nuclear receptor subfamily 0, group B, member 1 (NR0B1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	NR0B1 nuclear receptor subfamily 0, group B, member 1 [ Homo sapiens ]
<b>Official Symbol</b>	NR0B1
<b>Synonyms</b>	NR0B1; nuclear receptor subfamily 0, group B, member 1; AHC, dosage sensitive sex reversal , DSS; nuclear receptor subfamily 0 group B member 1; AHCH; DAX1; nuclear receptor DAX1; nuclear receptor DAX-1; nuclear hormone receptor; DSS-AHC critical region on the X chromosome protein 1; AHC; AHX; DSS; GTD; HHG; DAX-1; NROB1; SRXY2;
<b>Gene ID</b>	190
<b>mRNA Refseq</b>	NM_000475
<b>Protein Refseq</b>	NP_000466
<b>MIM</b>	300473
<b>UniProt ID</b>	P51843
<b>Chromosome Location</b>	Xp21.3
<b>Pathway</b>	Androgen Receptor Signaling Pathway, organism-specific biosystem; Gene Expression, organism-specific biosystem; Generic Transcription Pathway, organism-specific biosystem; Nuclear Receptor transcription pathway, organism-specific biosystem; Nuclear Receptors, organism-specific biosystem; Regulation of Androgen

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

receptor activity, organism-specific biosystem;

**Function**

AF-2 domain binding; DNA binding; DNA hairpin binding; RNA binding; double-stranded DNA binding; ligand-activated sequence-specific DNA binding RNA polymerase II transcription factor activity; ligand-activated sequence-specific DNA binding RNA polymerase II transcription factor activity; protein binding; protein domain specific binding; protein homodimerization activity; receptor activity; sequence-specific DNA binding; steroid hormone receptor activity; steroid hormone receptor binding; transcription corepressor activity; transcription factor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA