

Active Recombinant Human NR1H3, 1-182aa, GST-tagged

Cat. No. NR1H3-1058H Lot. No. (See product label)

SPECIFICATION

Species Human

Source E.coli

Description

Liver X Receptors (LXRs) are nuclear receptors that regulate the metabolism of cholesterol and bile acids. There are two subtypes of LXRs, LXR α and LXR β . LXR α is preferentially expressed in liver, small intestine, kidney and spleen. In contrast, LXR β expression is ubiquitous. The genomic structure and the promoter regions of the two LXR genes contain specific regulatory sites, which suggest that LXRs may have physiological roles in the immune system. Like other nuclear receptors, LXRs heterodimerize with Retinoid X Receptor (RXR) for function. LXRs are activated by naturally occurring oxysterols and regulate the expression of target genes, including ATP binding cassette transporter 1 (ABC1), ATP binding cassette transporter 8 (ABC8) and cholesterol ester transfer protein (CETP). LXR is thought to play a major role in the control of cholesterol catabolism by stimulating the expression of cholesterol 7 α -hydroxylase (CYP7A1), the rate limiting enzyme of bile acid synthesis. Recombinant GST-LXR α N182 is isolated from an *E. coli* strain that carries the coding sequence of the human LXR from amino acids 1-182 under the control of a T7 promoter.

Application GST-LXR α -N182 has been applied in DNA and protein-protein interaction assays.

Usage For in vitro use only.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	Liquid. Supplied in 20 mM Tris-HCl pH 8.0, 100 mM KCl, 0.2 mM EDTA, 1 mM DTT and 20% glycerol.
Activity	20 ng are sufficient for a gel-mobility shift assay and 100 ng are sufficient for a protein-protein interaction assay.
Purity	> 95% by SDS-PAGE.
Storage	Quality guaranteed for 12 months, Store at -80°C. Avoid freeze / thaw cycles.

GENE INFORMATION

Gene Name	NR1H3 nuclear receptor subfamily 1, group H, member 3 [Homo sapiens]
Synonyms	NR1H3; nuclear receptor subfamily 1, group H, member 3; LXRA; LXR-a; RLD-1; OTTHUMP00000198038; liver X receptor alpha; liver X receptor, alpha; liver X receptor-alpha; TLS/CHOP; Nuclear receptor subfamily 1 group H member 3; Nuclear orphan receptor LXR-alpha
Gene ID	10062
mRNA Refseq	NM_001130101
Protein Refseq	NP_001123573
MIM	602423
UniProt ID	Q13133
Chromosome Location	11p11.2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA


Pathway PPAR signaling pathway; Gene Expression; Metabolism of lipids and lipoproteins

Function


metal ion binding; protein binding; sequence-specific DNA binding; thyroid hormone receptor activity; transcription coactivator activity; zinc ion binding; steroid hormone receptor activity

PDB rendering based
on 1uhl.



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA