

Recombinant Human NR2C1 293 Cell Lysate

Cat. No. NR2C1-3714HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for nuclear receptor subfamily 2, group C, member 1 (NR2C1), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	NR2C1 nuclear receptor subfamily 2, group C, member 1 [Homo sapiens]
Official Symbol	NR2C1
Synonyms	NR2C1; nuclear receptor subfamily 2, group C, member 1; TR2; nuclear receptor subfamily 2 group C member 1; TR2 11; orphan nuclear receptor TR2; TR2 nuclear hormone receptor; nuclear receptor subfamily 2, group C isoform;
Gene ID	7181
mRNA Refseq	NM_001032287
Protein Refseq	NP_001027458
MIM	601529
UniProt ID	P13056
Chromosome Location	12q22
Pathway	Gene Expression, organism-specific biosystem; Generic Transcription Pathway, organism-specific biosystem; Nuclear Receptor transcription pathway, organism-specific biosystem; Regulation of Androgen receptor activity, organism-specific biosystem; Signaling events mediated by HDAC Class I, organism-specific biosystem;
Function	DNA binding; histone deacetylase binding; metal ion binding; protein binding; protein

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



homodimerization activity; receptor activity; sequence-specific DNA binding;
sequence-specific DNA binding transcription factor activity; steroid hormone receptor
activity; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA