

Recombinant Human NRG1 cell lysate

Cat. No. NRG1-1592HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human NRG1 (aa 177-241) derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Protein Length	177-241 a.a.
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[NRG1 neuregulin 1 \[Homo sapiens \]](#)

Official Symbol

NRG1

Synonyms

NRG1; neuregulin 1; HGL; pro-neuregulin-1, membrane-bound isoform; GGF; HRG; NDF; MSTP131; pro-NRG1; glial growth factor; neu differentiation factor; neuregulin 1 type IV beta 3; neuregulin 1 type IV beta 1a; sensory and motor neuron derived factor; heregulin, alpha (45kD, ERBB2 p185-activator); ARIA; GGF2; HRG1; HRGA; SMDF; MST131;

Gene ID

[3084](#)

mRNA Refseq

[NM_001159995](#)

Protein Refseq

[NP_001153467](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	142445
UniProt ID	Q02297
Chromosome Location	8p21-p12
Pathway	Downregulation of ERBB2:ERBB3 signaling, organism-specific biosystem; ErbB signaling pathway, organism-specific biosystem; ErbB signaling pathway, organism-specific biosystem; ErbB signaling pathway, conserved biosystem; GRB2 events in ERBB2 signaling, organism-specific biosystem; GRB7 events in ERBB2 signaling, organism-specific biosystem; Glypican 1 network, organism-specific biosystem;
Function	ErbB-3 class receptor binding; ErbB-3 class receptor binding; cytokine activity; growth factor activity; growth factor activity; protein binding; protein tyrosine kinase activator activity; receptor binding; receptor tyrosine kinase binding; transcription cofactor activity; transmembrane receptor protein tyrosine kinase activator activity; transmembrane receptor protein tyrosine kinase activator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA