

## Recombinant Human NRG1 cell lysate

**Cat. No.** NRG1-1675HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Human NRG1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
<b>Species</b>	Human
<b>Source</b>	Human Cells
<b>Preparation method</b>	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
<b>Lysis buffer</b>	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
<b>Quality control Testing</b>	12.5% SDS-PAGE Stained with Coomassie Blue
<b>Recommended Usage</b>	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

**Stability**

Samples are stable for up to twelve months from date of receipt at -80°C

**Storage Buffer**

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

**Storage Instruction**

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

## GENE INFORMATION

**Gene Name**

[NRG1 neuregulin 1 \[ Homo sapiens \]](#)

**Official Symbol**

NRG1

**Synonyms**

NRG1; neuregulin 1; HGL; pro-neuregulin-1, membrane-bound isoform; GGF; HRG; NDF; MSTP131; pro-NRG1; glial growth factor; neu differentiation factor; neuregulin 1 type IV beta 3; neuregulin 1 type IV beta 1a; sensory and motor neuron derived factor; heregulin, alpha (45kD, ERBB2 p185-activator); ARIA; GGF2; HRG1; HRGA; SMDF; MST131;

**Gene ID**

[3084](#)

**mRNA Refseq**

[NM\\_001159995](#)

**Protein Refseq**

[NP\\_001153467](#)

**MIM**

[142445](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>UniProt ID</b>	Q02297
<b>Chromosome Location</b>	8p21-p12
<b>Pathway</b>	Downregulation of ERBB2:ERBB3 signaling, organism-specific biosystem; ErbB signaling pathway, organism-specific biosystem; ErbB signaling pathway, organism-specific biosystem; ErbB signaling pathway, conserved biosystem; GRB2 events in ERBB2 signaling, organism-specific biosystem; GRB7 events in ERBB2 signaling, organism-specific biosystem; Glypican 1 network, organism-specific biosystem;
<b>Function</b>	ErbB-3 class receptor binding; ErbB-3 class receptor binding; cytokine activity; growth factor activity; growth factor activity; protein binding; protein tyrosine kinase activator activity; receptor binding; receptor tyrosine kinase binding; transcription cofactor activity; transmembrane receptor protein tyrosine kinase activator activity; transmembrane receptor protein tyrosine kinase activator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA