

# Recombinant Human/Bovine/Porcine NRG1 Protein

**Cat. No.** NRG1-29H    **Lot. No.** (See product label)

## SPECIFICATION

**Product Overview**

Recombinant human/bovine/porcine NRG-1 protein without tag was expressed in E. coli.

Animal-free (AOF) and carrier protein-free.

Mass spectrometry: single species with expected mass.

Analytical reversed-phase: single sharp peak.

Recovery from stock vial: >95%

**Species**                      Human

**Source**                        E.coli

**Description**

The protein encoded by this gene is a membrane glycoprotein that mediates cell-cell signaling and plays a critical role in the growth and development of multiple organ systems. An extraordinary variety of different isoforms are produced from this gene through alternative promoter usage and splicing. These isoforms are expressed in a tissue-specific manner and differ significantly in their structure, and are classified as types I, II, III, IV, V and VI. Dysregulation of this gene has been linked to diseases such as cancer, schizophrenia, and bipolar disorder (BPD).

**Molecular Mass**            7.5 kDa

**Endotoxin**                    <0.005 EU/μg protein (below level of detection)

**Applications**

Maintenance of pluripotency in human iPSC; cell culture media optimisation and replacement of insulin in defined media

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Storage** Resuspend in water at >100 µg/mL, prepare single use aliquots, add carrier protein if desired and store frozen at -20 or -80 centigrade

**Storage Buffer** Lyophilized from acetonitrile, TFA

## GENE INFORMATION

**Gene Name** [NRG1 neuregulin 1 \[ Homo sapiens \(human\) \]](#)

**Official Symbol** [NRG1](#)

**Synonyms** NRG1; neuregulin 1; GGF; HGL; HRG; NDF; ARIA; GGF2; HRG1; HRGA; SMDF; MST131; MSTP131; NRG1-IT2; pro-neuregulin-1, membrane-bound isoform; acetylcholine receptor-inducing activity; glial growth factor 2; heregulin, alpha (45kD, ERBB2 p185-activator); neu differentiation factor; pro-NRG1; sensory and motor neuron derived factor

**Gene ID** [3084](#)

**mRNA Refseq** [NM\\_013960](#)

**Protein Refseq** [NP\\_039254](#)

**MIM** [142445](#)


**UniProt ID** [Q02297](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

### Bioactivity



NRG-1 activity is determined using the serum response element luciferase reporter assay in transfected MCF-7 cells. EC<sub>50</sub> = 0.28 ng/mL (37.9 pM). Cells are treated in triplicate with a serial dilution of NRG-1 for 4 hours. Firefly luciferase activity is measured and normalized to the control Renilla luciferase activity.

### Purity




Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA



NRG-1 migrates as a single band at 7.5 kDa in non-reducing conditions and upon reduction (R). No contaminating protein bands are visible. Purified recombinant protein was resolved using 18% w/v SDS-PAGE in reduced and non-reduced conditions and stained with Coomassie Brilliant Blue R250.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA