

Recombinant Human NT5C2 293 Cell Lysate

Cat. No. NT5C2-3677HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for 5-nucleotidase, cytosolic II (NT5C2), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	NT5C2 5-nucleotidase, cytosolic II [Homo sapiens]
Official Symbol	NT5C2
Synonyms	NT5C2; 5-nucleotidase, cytosolic II; 5 nucleotidase (purine), cytosolic type B , NT5B; cytosolic purine 5-nucleotidase; cN II; GMP; PNT5; purine 5 nucleotidase; IMP-specific 5-NT; cytosolic 5-nucleotidase II; 5-nucleotidase (purine), cytosolic type B; NT5B; cN-II;
Gene ID	22978
mRNA Refseq	NM_001134373
Protein Refseq	NP_001127845
MIM	600417
UniProt ID	P49902
Chromosome Location	10q24.32
Pathway	Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of nucleotides, organism-specific biosystem; Nicotinate and nicotinamide metabolism, organism-specific biosystem; Nicotinate and nicotinamide metabolism, conserved biosystem; Purine catabolism, organism-specific biosystem; Purine metabolism, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

5-nucleotidase activity; hydrolase activity; metal ion binding; nucleotide binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA