

Recombinant Mouse NTRK2 cell lysate

Cat. No. NTRK2-1843MCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Mouse NTRK2 / TrkB derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[Ntrk2 neurotrophic tyrosine kinase, receptor, type 2 \[Mus musculus \]](#)

Official Symbol

NTRK2

Synonyms

NTRK2; neurotrophic tyrosine kinase, receptor, type 2; BDNF/NT-3 growth factors receptor; trk-B; GP145-TrkB/GP95-TrkB; trkB tyrosine kinase; Tkrb; trkB; AI848316; C030027L06Rik;

Gene ID

[18212](#)

mRNA Refseq

[NM_001025074](#)

Protein Refseq

[NP_001020245](#)

Pathway

Activation of TRKA receptors, organism-specific biosystem; MAPK signaling pathway, organism-specific biosystem; MAPK signaling pathway, conserved biosystem; NGF signalling via TRKA from the plasma membrane, organism-specific biosystem; NGF-independent TRKA activation, organism-specific biosystem; Neurotrophin signaling

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

pathway, organism-specific biosystem; Neurotrophin signaling pathway, conserved biosystem;

Function

ATP binding; brain-derived neurotrophic factor binding; brain-derived neurotrophic factor binding; brain-derived neurotrophic factor-activated receptor activity; brain-derived neurotrophic factor-activated receptor activity; kinase activity; neurotrophin receptor activity; neurotrophin-3 binding; neurotrophin-3 binding; neurotrophin-4/5 binding; nucleotide binding; protein binding; protein homodimerization activity; protein kinase activity; protein tyrosine kinase activity; receptor activity; transferase activity; transferase activity, transferring phosphorus-containing groups; transmembrane receptor protein tyrosine kinase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA