

Recombinant Human NUDT1 293 Cell Lysate

Cat. No. NUDT1-3656HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for nudix (nucleoside diphosphate linked moiety X)-type motif 1 (NUDT1), transcript variant 3B is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [NUDT1 nudix \(nucleoside diphosphate linked moiety X\)-type motif 1 \[Homo sapiens \]](#)

Official Symbol [NUDT1](#)

Synonyms

NUDT1; nudix (nucleoside diphosphate linked moiety X)-type motif 1; MTH1; 7,8-dihydro-8-oxoguanine triphosphatase; 7; 8 dihydro 8 oxoguanine triphosphatase; 8 oxo 7; 8 dihydrodeoxyguanosine triphosphatase; 8 dihydroguanosine triphosphatase; 8 oxo dGTPase; mutT human homolog 1; nucleoside diphosphate linked moiety X type motif 1; nudix motif 1; 8-oxo-dGTPase; 2-hydroxy-dATP diphosphatase; 8-oxo-7,8-dihydroguanosine triphosphatase; 8-oxo-7,8-dihydrodeoxyguanosine triphosphatase; nucleoside diphosphate-linked moiety X motif 1; nucleoside diphosphate-linked moiety X-type motif 1;

Gene ID [4521](#)

mRNA Refseq [NM_198952](#)

Protein Refseq [NP_945190](#)

MIM [600312](#)

UniProt ID [P36639](#)

Chromosome Location [7p22](#)

Pathway [oxidized GTP and dGTP detoxification, organism-specific biosystem; oxidized GTP](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



and dGTP detoxification, conserved biosystem;

Function

8-oxo-7,8-dihydrodeoxyguanosine triphosphate pyrophosphatase activity; 8-oxo-7,8-dihydroguanosine triphosphate pyrophosphatase activity; GTPase activity; hydrolase activity; metal ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA