

Recombinant Human OSM protein

Cat. No. OSM-25H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human OSM protein 227 a.a.) was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	227
Description	<p>Oncostatin M (OSM) is a multifunctional cytokine that belongs to the Interleukin-6 subfamily. Among the family members, OSM is most closely related to leukemia inhibitory factor (LIF) and it in fact utilizes the LIF receptor in addition to its specific receptor in the human. A biologically active OSM receptor has been previously described that consists of a heterodimer of leukemia inhibitory factor receptor (LIFR) and gp130. OSM is synthesized by stimulated T-cells and monocytes. The effects of OSM on endothelial cells suggest a pro-inflammatory role for OSM and endothelial cells possess a large number of OSM receptors. Recombinant murine OSM contains 181 amino acids and has a molecular mass of 20.4 kDa. It has approximately 48 % and 72 % amino acid sequence identity with human and rat OSM.</p>
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using human TF-1 cells is less than 2 ng/ml, corresponding to a specific activity of > 5.0 × 10 ⁵ IU/mg.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 25.8 kDa, a single non-glycosylated polypeptide chain containing 227 amino acids.
AA Sequence	AAIGSCSKEYRVLLGQLQKQTDLMQDTSRLDPYIRIQGLDVPKLRHCRERPGAFP SEETLRGLGRRGFLQTLNATLGCVLHRLADLEQRLPKAQDLERSGLNIEDLEKLQMA RPNILGLRNNIYCMAQLLDNSDTAEPTKAGRGASQPPTPTPASDAFQRKLEGCRFLH GYHRFMHSVGRVFSKWGESPNRSRRHSPHQALRKGVRRTSPSRKKGKRLMTRGQL PR
Endotoxin	Less than 1 EU/μg of rHuOSM as determined by LAL method.
Purity	>95% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	OSM
Official Symbol	OSM
Synonyms	OSM; oncostatin M; oncostatin-M; MGC20461;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID 5008


mRNA Refseq NM_020530

Protein Refseq NP_065391

MIM 165095

UniProt ID P13725

SDS-PAGE of OSM-
25H



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA