

Active Recombinant Human OSM Protein (228 aa)

Cat. No. OSM-3210 Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human Oncostatin M (rhOSM) produced in E. coli is a single non-glycosylated polypeptide chain containing 228 amino acids. A fully biologically active molecule, rhOSM has a molecular mass of 25.9 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques.
Species	Human
Source	E.coli
ProteinLength	228
Description	Oncostatin M (OSM) is a multifunctional cytokine, and belongs to Interleukin-6 (IL-6) subfamily, including IL-11, leukemia inhibitory factor (LIF), ciliary neurotropic factor, cardiotrophin-1, and novel neurotrophin-1. In vivo, OSM is secreted from activated T cells, monocytes, neutrophils, and endothelial cells. OSM is related to LIF, and share a receptor with LIF in human. Human OSM can bind to gp130 and recruit OSM Receptor β or LIF Receptor β to form a ternary complex. OSM stimulates the growth of different types of cells, including megakaryocytes, fibroblasts, vascular endothelial cells, and T cells. On the other hand, OSM inhibits the proliferation of several cancer cell lines, such as solid tissue tumor cells, lung cancer cells, melanoma cells, and breast cancer cells.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	ED50 < 10 ng/mL, measured by a cell proliferation assay using TF-1 cells,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

corresponding to a specific activity of > 1 × 10⁵ units/mg.

Molecular Mass 25.9 kDa, observed by reducing SDS-PAGE.

AA Sequence
 MAAIGSCSKEYRVLLGQLQKQTDLMQDTSRLDPYIRIQGLDVPKLRHCRERPGAF
 PSEETLRGLGRRGFLQTLNATLGCVLHRLADLEQRLPKAQDLERSGLNIEDLEKLQM
 ARPNILGLRNNIYCMAQLLDNSDTAEPTKAGRGASQPPTPTPASDAFQRKLEGCRFL
 HGYHRFMHSVGRVFSKWGESPNRSRRHSPHQALRKGVRRTTRPSRKGKRLMTRG
 QLPR

Endotoxin < 0.2 EU/μg, determined by LAL method.

Purity > 95% by SDS-PAGE analysis.

Storage Lyophilized recombinant human Oncostatin M (rhOSM) remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, rhOSM should be stable up to 2 weeks at 4 centigrade or up to 3 months at -20 centigrade.

Storage Buffer Lyophilized after extensive dialysis against PBS.

Reconstitution Reconstituted in ddH₂O at 100 μg/mL.

GENE INFORMATION

Gene Name OSM oncostatin M [Homo sapiens]

Official Symbol OSM

Synonyms OSM; oncostatin M; oncostatin-M; MGC20461;

Gene ID 5008

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



mRNA Refseq	NM_020530
Protein Refseq	NP_065391
MIM	165095
UniProt ID	P13725

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA