

Recombinant Human P4HB cell lysate

Cat. No. P4HB-2764HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human P4HB / ERBA2L derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

P4HB prolyl 4-hydroxylase, beta polypeptide [Homo sapiens]

Official Symbol

P4HB

Synonyms

P4HB; prolyl 4-hydroxylase, beta polypeptide; ERBA2L, PO4DB, procollagen proline, 2 oxoglutarate 4 dioxygenase (proline 4 hydroxylase), beta polypeptide , procollagen proline, 2 oxoglutarate 4 dioxygenase (proline 4 hydroxylase), beta polypeptide (protein disulfide isomerase associated 1) , procollagen proline, 2 oxoglutarate 4 dioxygenase (proline 4 hydroxylase), beta polypeptide (protein disulfide isomerase; thyroid hormone binding protein p55); protein disulfide-isomerase; collagen prolyl 4 hydroxylase beta; DSI; GIT; P4Hbeta; PDI; PDIA1; PO4HB; PROHB; protein disulfide isomerase family A; member 1; protein disulfide isomerase associated 1; p55; protocollagen hydroxylase; prolyl 4-hydroxylase subunit beta; collagen prolyl 4-hydroxylase beta; thyroid hormone-binding protein p55; glutathione-insulin transhydrogenase; cellular thyroid hormone-binding protein; protein disulfide isomerase-associated 1; protein disulfide isomerase/oxidoreductase; protein disulfide isomerase family A, member 1; procollagen-proline, 2-oxoglutarate 4-dioxygenase

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

(proline 4-hydroxylase), beta polypeptide; PHDB; PO4DB; ERBA2L;

Gene ID [5034](#)

mRNA Refseq [NM_000918](#)

Protein Refseq [NP_000909](#)

MIM [176790](#)

UniProt ID [P07237](#)

Chromosome Location 17q25

Pathway

Chylomicron-mediated lipid transport, organism-specific biosystem; Lipid digestion, mobilization, and transport, organism-specific biosystem; Lipoprotein metabolism, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of lipids and lipoproteins, organism-specific biosystem; Protein processing in endoplasmic reticulum, organism-specific biosystem; Protein processing in endoplasmic reticulum, conserved biosystem;

Function

electron carrier activity; isomerase activity; procollagen-proline 4-dioxygenase activity; protein binding; protein disulfide isomerase activity; protein disulfide oxidoreductase activity; protein heterodimerization activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA