

Recombinant Human PARP1 cell lysate

Cat. No. PARP1-710HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human PARP1 / PARP-1 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

PARP1 poly (ADP-ribose) polymerase 1 [Homo sapiens]

Official Symbol

PARP1

Synonyms

PARP1; poly (ADP-ribose) polymerase 1; ADP ribosyltransferase (NAD+; poly (ADP ribose) polymerase) , ADPRT, poly (ADP ribose) polymerase family, member 1 , PPOL; poly [ADP-ribose] polymerase 1; PARP; poly(ADP-ribose) polymerase; poly(ADP-ribose) synthetase; poly[ADP-ribose] synthase 1; poly(ADP-ribosyl)transferase; ADP-ribosyltransferase NAD(+); NAD(+) ADP-ribosyltransferase 1; poly (ADP-ribose) polymerase family, member 1; ADP-ribosyltransferase (NAD+; poly (ADP-ribose) polymerase); PPOL; ADPRT; ADPRT1; PARP-1; ADPRT 1; pADPRT-1;

Gene ID

142

mRNA Refseq

NM_001618

Protein Refseq

NP_001609

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	173870
UniProt ID	P09874
Chromosome Location	1q41-q42
Pathway	BER complex, organism-specific biosystem; BER complex, conserved biosystem; Base excision repair, organism-specific biosystem; Base excision repair, conserved biosystem; Caspase cascade in apoptosis, organism-specific biosystem; FAS pathway and Stress induction of HSP regulation, organism-specific biosystem; Notch-mediated HES/HEY network, organism-specific biosystem;
Function	DNA binding; NAD binding; NAD+ ADP-ribosyltransferase activity; metal ion binding; protein N-terminus binding; protein binding; transcription factor binding; transferase activity, transferring glycosyl groups; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA