

Recombinant Human PCCA 293 Cell Lysate

Cat. No. PCCA-3400HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for propionyl Coenzyme A carboxylase, alpha polypeptide (PCCA), nuclear gene encoding mitochondrial protein, transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [PCCA propionyl CoA carboxylase, alpha polypeptide \[Homo sapiens \]](#)

Official Symbol PCCA

Synonyms PCCA; propionyl CoA carboxylase, alpha polypeptide; propionyl Coenzyme A carboxylase, alpha polypeptide; propionyl-CoA carboxylase alpha chain, mitochondrial; PCCase alpha subunit; pccA complementation group; propanoyl-CoA:carbon dioxide ligase alpha subunit;

Gene ID [5095](#)

mRNA Refseq [NM_000282](#)

Protein Refseq [NP_000273](#)

MIM [232000](#)

UniProt ID [P05165](#)

Chromosome Location 13q32

Pathway 2-oxobutanoate degradation I, organism-specific biosystem; Fatty acid, triacylglycerol, and ketone body metabolism, organism-specific biosystem; Glyoxylate and dicarboxylate metabolism, organism-specific biosystem; Glyoxylate and dicarboxylate metabolism, conserved biosystem; Metabolic pathways, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Metabolism, organism-specific biosystem; Metabolism of lipids and lipoproteins, organism-specific biosystem;

Function

ATP binding; biotin binding; biotin carboxylase activity; enzyme binding; ligase activity; metal ion binding; nucleotide binding; propionyl-CoA carboxylase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA