

Recombinant Proliferating Cell Nuclear Antigen, GST-tagged

Cat. No. PCNA-4488H Lot. No. (See product label)

SPECIFICATION

Product Overview	PCNA full length Human Recombinant protein produced in E.Coli is a single, non-glycosylated polypeptide chain having a molecular mass of 62kDa which includes GST-tag fusion protein.
Species	Human
Source	Human
Description	PCNA is located in the nucleus and is a cofactor of DNA polymerase delta. PCNA acts as a homotrimer and helps elevate the processivity of leading strand synthesis during DNA replication. In reaction to DNA damage, PCNA is ubiquitinated and takes part in the RAD6-dependent DNA repair pathway. 2 transcript variants encoding the same protein have been found for PCNA gene. Pseudogenes of this PCNA gene have been described on chromosome 4 and on the X chromosome. PCNA is expressed during late G1- phase, S-phase of mitosis and persists until the end of the M-phase because of its long biological half-life. PCNA is induced by UV irradiation.
Form	The PCNA solution contains 50mM Tris-Acetate pH-7.5 & 1mM EDTA & 20% glycerol.
Physical Appearance	Sterile Filtered colorless solution.
Storage	PCNA although stable 4°C for 4 weeks, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathways Base excision repair; Cell cycle; DNA replication; Mismatch repair; Nucleotide excision repair; Cell Cycle, Mitotic; DNA Repair; DNA Replication; Telomere Maintenance

Full Length Full L.

GENE INFORMATION

Gene Name [PCNA proliferating cell nuclear antigen \[Homo sapiens \]](#)

Official Symbol [PCNA](#)

Synonyms MGC8367; PCNA; Proliferating cell nuclear antigen; Cyclin; proliferating cell nuclear antigen; cyclin; DNA polymerase delta auxiliary protein

Gene ID [5111](#)

mRNA Refseq [NM_002592](#)

Protein Refseq [NP_002583](#)

MIM [176740](#)

UniProt ID [P12004](#)

Chromosome Location 20pter-p12


Function DNA binding; DNA polymerase processivity factor activity; MutLalpha complex binding; dinucleotide insertion or deletion binding; protein binding; purine-specific mismatch base pair DNA N-glycosylase activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

The assembled
human DNA clamp, a
trimer of the protein
PCNA.



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA