

## Recombinant Rat PDGFRB cell lysate

**Cat. No.** PDGFRB-1112RCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Rat PDGFRB / PDGFR-1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
<b>Species</b>	Rat
<b>Source</b>	Human Cells
<b>Preparation method</b>	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
<b>Lysis buffer</b>	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
<b>Quality control Testing</b>	12.5% SDS-PAGE Stained with Coomassie Blue
<b>Recommended Usage</b>	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

**Stability**

Samples are stable for up to twelve months from date of receipt at -80°C

**Storage Buffer**

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

**Storage Instruction**

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

## GENE INFORMATION

**Gene Name**

[Pdgfrb platelet derived growth factor receptor, beta polypeptide \[ Rattus norvegicus \]](#)

**Official Symbol**

PDGFRB

**Synonyms**

PDGFRB; platelet derived growth factor receptor, beta polypeptide; platelet-derived growth factor receptor beta; PDGFR-beta; PDGF-R-beta; CD140 antigen-like family member B; platelet-derived growth factor receptor 1; beta platelet-derived growth factor receptor; Platelet-derived growth factor receptor, beta; beta-type platelet-derived growth factor receptor; platelet-derived growth factor receptor beta variant 1; PDGFR-1;

**Gene ID**

[24629](#)

**mRNA Refseq**

[NM\\_031525](#)

**Protein Refseq**

[NP\\_113713](#)

**Pathway**

Calcium signaling pathway, organism-specific biosystem; Calcium signaling pathway,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


conserved biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Downstream signal transduction, organism-specific biosystem; Focal adhesion, organism-specific biosystem; Focal adhesion, conserved biosystem;

**Function**

ATP binding; kinase activity; nucleotide binding; phosphatidylinositol 3-kinase binding; platelet-derived growth factor beta-receptor activity; platelet-derived growth factor beta-receptor activity; platelet-derived growth factor binding; platelet-derived growth factor receptor binding; platelet-derived growth factor-activated receptor activity; platelet-derived growth factor-activated receptor activity; protein binding; protein kinase activity; protein tyrosine kinase activity; protein tyrosine kinase activity; protein tyrosine kinase activity; receptor activity; receptor binding; signal transducer activity; vascular endothelial growth factor binding; vascular endothelial growth factor-activated receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA