

Recombinant Human PDK3 293 Cell Lysate

Cat. No. PDK3-3328HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for pyruvate dehydrogenase kinase, isozyme 3 (PDK3), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	PDK3 pyruvate dehydrogenase kinase, isozyme 3 [Homo sapiens]
Official Symbol	PDK3
Synonyms	PDK3; pyruvate dehydrogenase kinase, isozyme 3; pyruvate dehydrogenase kinase, isoenzyme 3; pyruvate dehydrogenase, lipoamide, kinase isozyme 3, mitochondrial;
Gene ID	5165
mRNA Refseq	NM_001142386
Protein Refseq	NP_001135858
MIM	602526
UniProt ID	Q15120
Chromosome Location	Xp22.12
Pathway	Metabolism, organism-specific biosystem; Pyruvate metabolism, organism-specific biosystem; Pyruvate metabolism and Citric Acid (TCA) cycle, organism-specific biosystem; Regulation of pyruvate dehydrogenase (PDH) complex, organism-specific biosystem; TCA Cycle, organism-specific biosystem; The citric acid (TCA) cycle and respiratory electron transport, organism-specific biosystem;
Function	ATP binding; nucleotide binding; protein binding; protein kinase activity; pyruvate


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA



dehydrogenase (acetyl-transferring) kinase activity; transferase activity; two-component sensor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA