

Recombinant Human Phosphoinositide-3-kinase, Regulatory Subunit 2 (Beta)

Cat. No. PIK3R2-51H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human Phosphoinositide 3-kinase beta produced in Sf9 is a glycosylated protein having a molecular weight as follows: p85 α chain 83.5 kDa, p110 β chain 124.3 kDa.
Species	Human
Source	Sf9 Cells
Description	The PI3K β isoform can be activated by insulin via the insulin receptor to initiate a cascade of events that control cell growth and metabolism. The activation of PI3K β is mediated by the p85 regulatory subunit binding to tyrosine phosphorylated insulin receptor substrate (IRS) proteins (e.g. IRS-1 and IRS-2). It was also shown that PI3K β is involved in apoptosis in human colon carcinoma cells. Injection of neutralizing antibodies specific to p110 β in WiDr, HCT116 and CO 115 adenocarcinoma cells inhibited de novo DNA synthesis. PI3K β is the major PI3K isoform required for apoptotic cell and Fc-g receptor mediated phagocytosis shown for primary mouse macrophages and the Jurkat human leukemia T cell line. It was shown by several research groups that the catalytic subunit of PI3K β can be activated by G β subunits of G-protein coupled receptors.
Physical Appearance	Sterile filtered liquid formulation.
Purity	Greater than 90.0% as determined by SDS Page.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Formulation	0.5 mg/ml solution in 10mM Hepes, pH 7.5, 100mM NaCl, 2.5mM MgCl ₂ , and 50% glycerol.
Biological Activity	3 nmol/mg/min using phosphatidylinositol as the substrate.
Storage	PI3Kb although stable at 14°C for 1 week, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.
GENE INFORMATION	
Gene Name	PIK3R2 phosphoinositide-3-kinase, regulatory subunit 2 (beta) [Homo sapiens]
Synonyms	PIK3R2; phosphoinositide-3-kinase, regulatory subunit 2 (beta); p85; P85B; p85-BETA; phosphoinositide-3-kinase, regulatory subunit 2 (p85 beta); phosphoinositide-3-kinase, regulatory subunit, polypeptide 2 (p85 beta); phosphatidylinositol 3-kinase, regulatory subunit, polypeptide 2 (p85 beta); Phosphatidylinositol 3-kinase regulatory subunit beta; PI3-kinase p85 subunit beta; PtdIns-3-kinase p85-beta
Gene ID	5296
mRNA Refseq	NM_005027
Protein Refseq	NP_005018
MIM	603157
UniProt ID	O00459
Chromosome Location	19q13.2-q13.4

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

Acute myeloid leukemia; Apoptosis; B cell receptor signaling pathway; Chemokine signaling pathway; Chronic myeloid leukemia; Colorectal cancer; Endometrial cancer; ErbB signaling pathway; Fc epsilon RI signaling pathway; Fc gamma R-mediated phagocytosis; Focal adhesion; Glioma; Insulin signaling pathway; Jak-STAT signaling pathway; Leukocyte transendothelial migration; Melanoma; Natural killer cell mediated cytotoxicity; Neurotrophin signaling pathway; Non-small cell lung cancer; Pancreatic cancer; Pathways in cancer; Phosphatidylinositol signaling system; Prostate cancer; Regulation of actin cytoskeleton; Renal cell carcinoma; Small cell lung cancer; T cell receptor signaling pathway; Toll-like receptor signaling pathway; Type II diabetes mellitus; VEGF signaling pathway; mTOR signaling pathway; Hemostasis; Signaling by Insulin receptor; Signaling by PDGF; Signaling by Rho GTPases; Signaling in Immune system; Signalling by NGF

Function

1-phosphatidylinositol-3-kinase activity; phosphoinositide 3-kinase regulator activity; protein binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA