

## Recombinant Human PLA2G2A cell lysate

**Cat. No.** PLA2G2A-2184HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Human PLA2G2A / PLA2B derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
<b>Species</b>	Human
<b>Source</b>	Human Cells
<b>Preparation method</b>	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
<b>Lysis buffer</b>	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
<b>Quality control Testing</b>	12.5% SDS-PAGE Stained with Coomassie Blue
<b>Recommended Usage</b>	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

**Stability**

Samples are stable for up to twelve months from date of receipt at -80°C

**Storage Buffer**

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

**Storage Instruction**

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

## GENE INFORMATION

**Gene Name**

PLA2G2A phospholipase A2, group IIA (platelets, synovial fluid) [ Homo sapiens ]

**Official Symbol**

PLA2G2A

**Synonyms**

PLA2G2A; phospholipase A2, group IIA (platelets, synovial fluid); PLA2B, PLA2L; phospholipase A2, membrane associated; NPS-PLA2; GIIC sPLA2; group IIA phospholipase A2; phosphatidylcholine 2-acylhydrolase 2A; non-pancreatic secretory phospholipase A2; MOM1; PLA2; PLA2B; PLA2L; PLA2S; PLAS1; sPLA2;

**Gene ID**

5320

**mRNA Refseq**

NM\_000300

**Protein Refseq**

NP\_000291

**MIM**


172411

**UniProt ID**

P14555

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Chromosome Location</b>	1p35
<b>Pathway</b>	Arachidonic acid metabolism, organism-specific biosystem; Arachidonic acid metabolism, conserved biosystem; Eicosanoid Synthesis, organism-specific biosystem; Ether lipid metabolism, organism-specific biosystem; Ether lipid metabolism, conserved biosystem; Fat digestion and absorption, organism-specific biosystem; Fat digestion and absorption, conserved biosystem;
<b>Function</b>	calcium ion binding; calcium-dependent phospholipase A2 activity; hydrolase activity; phospholipase A2 activity; phospholipid binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA