

Active Recombinant Human PLAT protein, His-tagged

Cat. No. PLAT-051H Lot. No. (See product label)

SPECIFICATION

| | |
|-------------------------|---|
| Product Overview | Active Recombinant Human PLAT protein(P00750)(311~562 aa), fused with N-terminal His tag, was expressed in HEK293. |
| Species | Human |
| Source | HEK293 |
| ProteinLength | 311-562 aa |
| Form | Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. |
| Bio-activity | <p>Plasminogen activators are serine proteases that catalyze the activation of plasmin via a proteolytic cleavage of its zymogen form plasminogen. Plasmin is an important factor in fibrinolysis, the breakdown of fibrin polymers formed during blood clotting. There are two main plasminogen activators: urokinase (uPA) and tissue plasminogen activator (tPA). Tissue plasminogen activators (tPA) are used to treat medical conditions related to blood clotting including embolic or thrombotic stroke, myocardial infarction, and pulmonary embolism. The activity assay of tPA was measured by its ability to cleave a peptide substrate, N-carbobenzyloxy-Gly-Gly-Arg-7-amido-4-methylcoumarin (Z-GG R-AMC). The reaction was performed in 50 mM Tris, 0.01% Tween-20, pH 8.5 (Assay Buffer), initiated by addition 50 µL of various concentrations of tPA (diluted by Assay Buffer) to 50 µ of 200 uM Substrate. Read at excitation and emission wavelengths of 380 nm and 460 nm (top read), respectively, in kinetic mode for 5 minutes. The spe</p> |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

cific activity of recombinant human tPA is 10-60 pmol/min/g.

| | |
|-----------------------|--|
| Molecular Mass | 29.7kDa |
| Endotoxin | <1.0EU per 1µg (determined by the LAL method). |
| Purity | > 95% as determined by SDS-PAGE. |
| Storage | Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months. |
| Reconstitution | Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex. |

GENE INFORMATION

| | |
|------------------------|--|
| Gene Name | PLAT plasminogen activator, tissue [Homo sapiens] |
| Official Symbol | PLAT |
| Synonyms | PLAT; plasminogen activator, tissue; tissue-type plasminogen activator; alteplase; reteplase; t-plasminogen activator; plasminogen activator, tissue type; tissue plasminogen activator (t-PA); TPA; T-PA; DKFZp686I03148; |
| Gene ID | 5327 |
| mRNA Refseq | NM_000930 |
| Protein Refseq | NP_000921 |
| MIM | 173370 |


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID

P00750

SDS-PAGE Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA