

Recombinant Human PLAU cell lysate

Cat. No. PLAU-1734HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human Urokinase / PLAU (activated by trypsin) derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name [PLAU plasminogen activator, urokinase \[Homo sapiens \]](#)

Official Symbol [PLAU](#)

Synonyms [PLAU](#); plasminogen activator, urokinase; urokinase-type plasminogen activator; UPA; URK; U-plasminogen activator; plasminogen activator, urinary; ATF; QPD; u-PA; BDPLT5;

Gene ID [5328](#)

mRNA Refseq [NM_001145031](#)

Protein Refseq [NP_001138503](#)

MIM [191840](#)

UniProt ID [P00749](#)

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	10q24
Pathway	ATF-2 transcription factor network, organism-specific biosystem; Blood Clotting Cascade, organism-specific biosystem; Complement and Coagulation Cascades, organism-specific biosystem; Complement and coagulation cascades, organism-specific biosystem; Complement and coagulation cascades, conserved biosystem; DNA damage response (only ATM dependent), organism-specific biosystem; Dissolution of Fibrin Clot, organism-specific biosystem;
Function	peptidase activity; protein binding; serine-type endopeptidase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA