

## Recombinant Human POU5F1 293 Cell Lysate

**Cat. No.** POU5F1-2999HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	1-360 a.a.
<b>Description</b>	Antigen standard for POU class 5 homeobox 1 (POU5F1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Applications**

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

**GENE INFORMATION**
**Gene Name**

[POU5F1](#) POU class 5 homeobox 1 [ [Homo sapiens](#) ]

**Official Symbol**

[POU5F1](#)

**Synonyms**

[POU5F1](#); POU class 5 homeobox 1; OTF3, POU domain class 5, transcription factor 1; POU domain, class 5, transcription factor 1; [MGC22487](#); [OCT3](#); [Oct4](#); octamer-binding protein 3; octamer-binding protein 4; POU domain transcription factor [OCT4](#); octamer-binding transcription factor 3; octamer-binding transcription factor-3; POU-type homeodomain-containing DNA-binding protein; [OCT4](#); [OTF3](#); [OTF4](#); [OTF-3](#); [Oct-3](#); [Oct-4](#);

**Gene ID**

[5460](#)

**mRNA Refseq**

[NM\\_002701](#)

**Protein Refseq**

[NP\\_002692](#)

**MIM**

[164177](#)

**UniProt ID**

[Q01860](#)

**Chromosome Location**

6p21.33

**Pathway**

HIF-2-alpha transcription factor network, organism-specific biosystem; Wnt Signaling

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway and Pluripotency, organism-specific biosystem;

**Function**

DNA binding; miRNA binding; protein binding; sequence-specific DNA binding; sequence-specific DNA binding RNA polymerase II transcription factor activity; sequence-specific DNA binding transcription factor activity; transcription factor binding; transcription regulatory region DNA binding; ubiquitin protein ligase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA