

Active Recombinant Human PPARA, LB Domain, 170-468aa, His-tagged

Cat. No. PPARA-1085H Lot. No. (See product label)

SPECIFICATION

Species Human

Source E.coli

ProteinLength 170-468 a.a.

Description

Recombinant His tagged PPAR α -LBD(33 kDa) is isolated from an E.coli strain that carries the coding sequence of the human PPAR α under the control of a T7 promoter. There is evidence that a group of closely related nuclear receptors, called Peroxisome Proliferator-activated Receptors (PPARs), may be involved in chronic diseases such as diabetes, obesity, atherosclerosis and cancer. The PPARs were first cloned as the nuclear receptors that mediate the effects of synthetic compounds called peroxisome proliferators on gene transcription. It soon became clear that eicosanoids and fatty acids can also regulate gene transcription through PPARs. They bind a specific element in the promoter region of target genes only as a heterodimer with the receptor for 9-cis retinoic acid, RXR (retinoid X receptor). Binding of the ligand of either receptor can activate the complex, but binding of both ligands simultaneously is more potent. Three PPAR isotypes have been identified: α , β (also called NUC1) and γ . PPAR α is expressed most in brown adipose tissue and liver, then kidney, heart and skeletal muscle. PPAR γ is mainly expressed in adipose tissue, and to a lesser extent in colon, the immune system and the retina. PPAR β is found in many tissues but the highest expression is in the gut, kidney and heart. The target genes of PPAR α are a relatively homogenous group of genes that participate in

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

aspects of lipid catabolism such as fatty acid uptake through membranes, fatty acid binding in cells, fatty acid oxidation (in microsomes, peroxisomes and mitochondria) and lipoprotein assembly and transport.

Form

Liquid. Supplied in 20 mM Tris-HCl pH 8.0, 20% glycerol, 100 mM KCl, 0.2 mM EDTA and 1 mM DTT.

Purity

> 95% by SDS-PAGE.

Application

PPAR α has been applied in DNA and protein-protein interaction assays.

Activity

20-100 ng are sufficient for a ligand binding assay and 100 ng are sufficient for a protein-protein interaction assay.

Usage

For in vitro use only.

Storage

Quality guaranteed for 12 months store at -80°C. Avoid freeze / thaw cycles.

GENE INFORMATION

Gene Name

PPARA peroxisome proliferator-activated receptor alpha [Homo sapiens]

Synonyms

PPARA; peroxisome proliferator-activated receptor alpha; PPAR; NR1C1; hPPAR; MGC2237; MGC2452; PPARalpha; peroxisome proliferative activated receptor, alpha; OTTHUMP00000197740; OTTHUMP00000197741; PPAR-alpha; Nuclear receptor subfamily 1 group C member 1

Gene ID

5465

mRNA Refseq

NM_001001928


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_001001928
MIM	170998
UniProt ID	Q07869
Chromosome Location	22q12-q13.1
Pathway	Adipocytokine signaling pathway; PAR signaling pathway; Metabolism of lipids and lipoproteins
Function	drug binding; ligand-regulated transcription factor activity; lipid binding; metal ion binding; protein domain specific binding; sequence-specific DNA binding; steroid hormone receptor activity; transcription activator activity; transcription factor activity; zinc ion binding

PDB rendering based on 1i7g.



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA