

Active Recombinant Human PPIA, N-His tagged

Cat. No. PPIA-26334TH Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human PPIA, was expressed in E.coli and purified by using conventional chromatography techniques.
Species	Human
Source	E.coli
ProteinLength	1-165 aa
Description	This gene encodes a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. The encoded protein is a cyclosporin binding-protein and may play a role in cyclosporin A-mediated immunosuppression. The protein can also interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions. Multiple pseudogenes that map to different chromosomes have been reported.
Form	In 20mM Tris pH 8.0, .20mM NaCl, 0.5mM DTT, 10% glycerol
Bio-activity	Specific activity is > 270 nmoles/min/mg, and is defined as the amount of enzyme that cleaves 1umole of suc-AAFP-pNA per minute at 25C in Tris-Hcl pH8.0 using chymotrypsin.
Molecular Mass	20kDa (185aa), confirmed by MALDI-TOF.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	MGSSHHHHHH SSSLVPRGSH MVNPTVFFDI AVDGEPLGRV SFELFADKVP KTAENFRALS TGEKGFYKYG SCFHRIIPGF MCQGGDFTRH NGTGGKSIYG EKFEDENFIL KHTGPGILSM ANAGPNTNGS QFFICTAKTE WLDGKHVVFG KVKEGMNIVE AMERFGSRNG KTSKKITIAD CGQLE
Endotoxin	< 1.0 EU per 1 microgram of protein (determined by LAL method)
Purity	> 95% by SDS - PAGE
Applications	SDS-PAGE
Storage	Can be stored at +4C short term (1-2 weeks). For long term storage, aliquot and store at -20C or -70C. Avoid repeated freezing and thawing cycles.
Concentration	1 mg/ml (determined by Bradford assay)

GENE INFORMATION

Gene Name	PPIA peptidylprolyl isomerase A (cyclophilin A) [Homo sapiens]
Official Symbol	PPIA
Synonyms	PPIA; peptidylprolyl isomerase A (cyclophilin A); peptidyl-prolyl cis-trans isomerase A; CYPA; CYPH; HEL-S-69p; PPIase A; rotamase A; cyclophilin; T cell cyclophilin; cyclosporin A-binding protein; epididymis secretory sperm binding protein Li 69p; NP_066953.1; EC 5.2.1.8
Gene ID	5478
mRNA Refseq	NM_021130

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_066953
MIM	123840
UniProt ID	P62937
Chromosome Location	7p13
Pathway	APOBEC3G mediated resistance to HIV-1 infection; Assembly Of The HIV Virion; Basigin interactions
Function	peptide binding; peptidyl-prolyl cis-trans isomerase activity; poly(A) RNA binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA