

Recombinant Full Length Human PROCR Protein, His-tagged

Cat. No. PROCR-418H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human PROCR protein with His tag was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	1-238 a.a.
Description	The protein encoded by this gene is a receptor for activated protein C, a serine protease activated by and involved in the blood coagulation pathway. The encoded protein is an N-glycosylated type I membrane protein that enhances the activation of protein C. Mutations in this gene have been associated with venous thromboembolism and myocardial infarction, as well as with late fetal loss during pregnancy. The encoded protein may also play a role in malarial infection and has been associated with cancer.
Form	Lyophilized
Molecular Mass	22.8 kDa
AA Sequence	MLTTLPIILLSGWAFCSQDASDGLQRLHMLQISYFRDPYHVWYQGNASLGGHLTH VLEGPDTNTTIIQLQPLQEPESWARTQSGLQSYLLQFHGLVRLVHQERTLAFPLTIRC FLGCELPPEGSRAHVFFEVAVNGSSSFVSFRPERALWQADTQVTSGVVFTLQQLNA YNRTRYELREFLEDTCVQYVQKHISAENTKGSQTSRSYTSVLGVLVGSFIIAGVAVG IFLCTGGRRC

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	> 98%
Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Concentration	1 mg/mL
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H2O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.

GENE INFORMATION

Gene Name	PROCR protein C receptor, endothelial [Homo sapiens (human)]
Official Symbol	PROCR
Synonyms	PROCR; protein C receptor, endothelial; endothelial protein C receptor; CCD41; CD201; EPCR; APC receptor; centrocyclin; CD201 antigen; activated protein C receptor; endothelial cell protein C receptor; cell cycle, centrosome-associated protein; CCCA; bA42O4.2; MGC23024;
Gene ID	10544

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



mRNA Refseq	NM_006404
Protein Refseq	NP_006395
MIM	600646
UniProt ID	Q9UNN8

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA