

Recombinant Human PSMA1 293 Cell Lysate

Cat. No. PSMA1-2780HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for proteasome (prosome, macropain) subunit, alpha type, 1 (PSMA1), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [PSMA1 proteasome \(prosome, macropain\) subunit, alpha type, 1 \[Homo sapiens \]](#)

Official Symbol [PSMA1](#)

Synonyms [PSMA1](#); proteasome (prosome, macropain) subunit, alpha type, 1; proteasome subunit alpha type-1; HC2; MGC1667; MGC14542; MGC14575; MGC14751; MGC21459; MGC22853; MGC23915; NU; PROS30; PROS-30; protein P30-33K; proteasome nu chain; macropain subunit C2; macropain subunit nu; proteasome subunit nu; 30 kDa prosomal protein; proteasome component C2; proteasome subunit, alpha-type, 1; multicatalytic endopeptidase complex subunit C2;

Gene ID [5682](#)

mRNA Refseq [NM_001143937](#)

Protein Refseq [NP_001137409](#)

MIM [602854](#)

UniProt ID [P25786](#)

Chromosome Location [11p15.1](#)

Pathway [APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem](#); [APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem](#); [APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem](#);

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem;

Function

RNA binding; peptidase activity; protein binding; threonine-type endopeptidase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA