

Active Recombinant Sheep PTGS2 Protein

Cat. No. PTGS2-046S Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Sheep PTGS2 Protein (24-599) without tag was expressed in Sf21 cells using the Baculovirus system.
Species	Sheep
Source	Insect Cells
ProteinLength	24-599
Description	<p>Dual cyclooxygenase and peroxidase in the biosynthesis pathway of prostanoids, a class of C20 oxylipins mainly derived from arachidonate, with a particular role in the inflammatory response (PubMed:10438452). The cyclooxygenase activity oxygenates arachidonate (AA, C20:4(n-6)) to the hydroperoxy endoperoxide prostaglandin G2 (PGG2), and the peroxidase activity reduces PGG2 to the hydroxy endoperoxide PGH2, the precursor of all 2-series prostaglandins and thromboxanes (PubMed:10438452). This complex transformation is initiated by abstraction of hydrogen at carbon 13 (with S-stereochemistry), followed by insertion of molecular O2 to form the endoperoxide bridge between carbon 9 and 11 that defines prostaglandins. The insertion of a second molecule of O2 (bis-oxygenase activity) yields a hydroperoxy group in PGG2 that is then reduced to PGH2 by two electrons (By similarity). Similarly catalyzes successive cyclooxygenation and peroxidation of dihomogamma-linoleate (DGLA, C20:3(n-6)) and eicosapentaenoate (EPA, C20:5(n-3)) to corresponding PGH1 and PGH3, the precursors of 1- and 3-series prostaglandins (By similarity). In an alternative pathway of prostanoid biosynthesis,</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

converts 2-arachidonoyl lysophospholipids to prostanoid lysophospholipids, which are then hydrolyzed by intracellular phospholipases to release free prostanoids (By similarity). Metabolizes 2-arachidonoyl glycerol yielding the glyceryl ester of PGH₂, a process that can contribute to pain response. Generates lipid mediators from n-3 and n-6 polyunsaturated fatty acids (PUFAs) via a lipoxygenase-type mechanism. Oxygenates PUFAs to hydroperoxy compounds and then reduces them to corresponding alcohols (By similarity). Plays a role in the generation of resolution phase interaction products (resolvins) during both sterile and infectious inflammation. Metabolizes docosahexaenoate (DHA, C₂₂:6(n-3)) to 17R-HDHA, a precursor of the D-series resolvins (RvDs) (By similarity). As a component of the biosynthetic pathway of E-series resolvins (RvEs), converts eicosapentaenoate (EPA, C₂₀:5(n-3)) primarily to 18S-HEPE that is further metabolized by ALOX₅ and LTA₄H to generate 18S-RvE1 and 18S-RvE2 (By similarity). In vascular endothelial cells, converts docosapentaenoate (DPA, C₂₂:5(n-3)) to 13R-HDPA, a precursor for 13-series resolvins (RvTs) shown to activate macrophage phagocytosis during bacterial infection (By similarity). In activated leukocytes, contributes to oxygenation of hydroxyeicosatetraenoates (HETE) to diHETES (5,15-diHETE and 5,11-diHETE) (By similarity). During neuroinflammation, plays a role in neuronal secretion of specialized preresolving mediators (SPMs) 15R-lipoxin A₄ that regulates phagocytic microglia (By similarity).

Bio-activity Specific Activity: > 20000 U/mg

Molecular Mass ~70 kDa

Purity ≥80% as determined by SDS-PAGE

Unit Definition One unit is defined as the amount of enzyme required to consume 1 nmol of oxygen per minute at 37 centigrade in 100 mM Tris, pH 8.0, containing 100 μM arachidonate, 5 mM EDTA, 2 mM phenol, and 1 μM hematin.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Stability	≥ 1 year
Storage	At -80 centigrade.
GENE INFORMATION	
Gene Name	PTGS2 prostaglandin-endoperoxide synthase 2 [<i>Ovis aries</i> (sheep)]
Official Symbol	PTGS2
Synonyms	PTGS2; prostaglandin-endoperoxide synthase 2; PGHS-2; prostaglandin G/H synthase 2; COX-2; PGH synthase 2; PHS II; cyclooxygenase, prostaglandin endoperoxide H synthase-2; cyclooxygenase-2; prostaglandin H synthase-2; prostaglandin H2 synthase 2; prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase); EC 1.14.99.1
Gene ID	443460
mRNA Refseq	NM_001009432
Protein Refseq	NP_001009432
UniProt ID	P79208

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**SDS-PAGE analysis
of PTGS2**



Lane 1: MW Markers
Lane 2: PTGS2 (2 µg)
Lane 3: PTGS2 (5 µg)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA