

## Recombinant Human PTGS2 cell lysate

**Cat. No.** PTGS2-643HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Human PTGS2 / COX2 / PGHS-2 derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
<b>Species</b>	Human
<b>Source</b>	Insect Cells
<b>Preparation method</b>	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
<b>Lysis buffer</b>	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
<b>Quality control Testing</b>	12.5% SDS-PAGE Stained with Coomassie Blue
<b>Recommended Usage</b>	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

**Stability**

Samples are stable for up to twelve months from date of receipt at -80°C

**Storage Buffer**

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

**Storage Instruction**

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

**GENE INFORMATION**

**Gene Name**

PTGS2 prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase) [ Homo sapiens ]

**Official Symbol**

PTGS2

**Synonyms**

PTGS2; prostaglandin-endoperoxide synthase 2 (prostaglandin G/H synthase and cyclooxygenase); prostaglandin G/H synthase 2; COX2; PHS II; PGH synthase 2; cyclooxygenase 2b; prostaglandin H2 synthase 2; COX-2; PHS-2; PGG/HS; PGHS-2; hCox-2; GRIPGHS;

**Gene ID**

5743

**mRNA Refseq**

NM\_000963

**Protein Refseq**

NP\_000954

**MIM**

600262

Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

<b>UniProt ID</b>	P35354
<b>Chromosome Location</b>	1q25.2-q25.3
<b>Pathway</b>	Arachidonic acid metabolism, organism-specific biosystem; Arachidonic acid metabolism, conserved biosystem; Biological oxidations, organism-specific biosystem; C-MYB transcription factor network, organism-specific biosystem; COX reactions, organism-specific biosystem; Calcineurin-regulated NFAT-dependent transcription in lymphocytes, organism-specific biosystem; Calcium signaling in the CD4+ TCR pathway, organism-specific biosystem;
<b>Function</b>	enzyme binding; heme binding; heme binding; lipid binding; metal ion binding; oxidoreductase activity; oxidoreductase activity, acting on single donors with incorporation of molecular oxygen, incorporation of two atoms of oxygen; peroxidase activity; peroxidase activity; peroxidase activity; prostaglandin-endoperoxide synthase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA