

# Active Recombinant Human PVR protein, Fc-tagged, Biotinylated

**Cat. No.** PVR-051H    **Lot. No.** (See product label)

## SPECIFICATION

### Product Overview

Biotinylated Recombinant Human PVR(Asp28-Asn343) protein, fused to Fc tag at the C-terminus, was expressed in HEK293 cells .

### Species

Human

### Source

HEK293

### ProteinLength

Asp28-Asn343

### Description

CD155, also known as PVR (poliovirus receptor), Nectin-5 (nectin-like molecule-5) and, in rodents, TAGE4 (tumor-associated glycoprotein E4), is a 70 kDa type I transmembrane glycoprotein in the nectin-related family of adhesion proteins within the immunoglobulin superfamily (1, 2). CD155 binds other molecules including Vitronectin, Nectin-3, DNAM-1/CD226, CD96, and TIGIT but does not bind homotypically (3). Mature human CD155 consists of a 323 amino acid (aa) extracellular domain (ECD) with one N-terminal V-type and two C2-type Ig-like domains, a 24 aa transmembrane segment, and a 50 aa cytoplasmic tail. Within the ECD, human CD155 shares 45% aa sequence identity with mouse and rat CD155, and 52% with human Nectin-2. The V-type domain of CD155 mediates all binding, including to polio virus (1), and alternative splicing within this domain in humans can modulate ligand binding (4). Human CD155 can also be spliced to generate secreted isoforms (5). CD155 is up-regulated on endothelial cells by IFN-gamma and is highly expressed on immature thymocytes, lymph node dendritic cells, and tumor cells of

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

epithelial and neuronal origin (1, 2, 6-9). It is preferentially expressed on Th17 cells compared to Th2 cells (10), and its activation promotes the development of Th1 responses (11). On migrating cells, CD155 is concentrated at the leading edge, where it binds basement membrane Vitronectin, recruits Nectin-3-expressing cells, and cooperates with PDGF and Integrin alpha v beta 3 to promote cell migration (1, 3, 12). Enhanced CD155 expression in tumor cells contributes to loss of contact inhibition and increased migration but also allows tumor cell recognition and killing by DNAM-1 or CD96 expressing NK cells (1, 7, 13). Binding of monocyte DNAM-1 to endothelial cell CD155 promotes transendothelial migration (8). The expression of CD155 on mouse CD8+ thymocytes prevents their premature exit from the thymus (14). Within intestinal Peyer's patches, follicular dendritic cell CD155 activates follicular helper T cells via DNAM-1 or CD96 binding (7-9, 15). CD155 also binds the inhibitory ligand TIGIT on NK and some mature T cells, antagonizing DNAM-1 effects (7, 15, 16).

<b>Predicted N Terminal</b>	Asp28
<b>Form</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.
<b>Bio-activity</b>	Measured by its binding ability in a functional ELISA. When Recombinant Human CD96v2 Fc Chimera is coated at 1 ug/mL, 100 µL/well, the concentration of Biotinylated Recombinant Human PVR Fc Chimera that produces 50% of the optimal binding response is 8-48 ng/mL.
<b>Endotoxin</b>	<0.10 EU per 1 µg of the protein by the LAL method.
<b>Purity</b>	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
<b>Applications</b>	Bioactivity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Storage** Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.

**Reconstitution** Reconstitute at 500 µg/mL in PBS.

**Conjugation** Biotin

## GENE INFORMATION

**Gene Name** [PVR poliovirus receptor \[ Homo sapiens \]](#)

**Official Symbol** [PVR](#)

**Synonyms** PVR; poliovirus receptor; PVS; CD155; HVED; Necl 5; NECL5; nectin like 5; Tage4; nectin-like 5; nectin-like protein 5; TAGE4; Necl-5; FLJ25946;

**Gene ID** [5817](#)

**mRNA Refseq** [NM\\_001135768](#)

**Protein Refseq** [NP\\_001129240](#)

**MIM** [173850](#)

**UniProt ID** [P15151](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA