

Active Recombinant Human PVRIG protein, Fc-Avi-tagged, Biotinylated

Cat. No. PVRIG-523HB **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Biotinylated Recombinant Human PVRIG protein(Thr 41 - Asp 171), fused with Fc and Avi tag, was expressed in HEK293.

Species

Human

Source

HEK293

ProteinLength

Thr 41 - Asp 171

Form

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4, 10% trehalose.

Bio-activity

Immobilized Biotinylated Human PVRIG, Fc,Avitag at 5 µg/mL (100 µL/well) on streptavidin precoated (0.5 µg/well) plate can bind Human Nectin-2, Mouse IgG2a Fc Tag with a linear range of 0.1-3 ng/mL. Immobilized Biotinylated Human PVRIG, Fc,Avitag at 5 µg/mL (100 µL/well) on streptavidin precoated (0.5 µg/well) plate can bind Human Nectin-2, His Tag with a linear range of 0.02-0.313 µg/mL. Immobilized Human Nectin-2, His Tag at 10 µg/mL (100 µL/well) can bind Biotinylated Human PVRIG, Fc,Avitag with a linear range of 0.005-0.313 µg/mL. Immobilized Human Nectin-2, Fc Tag at 10 µg/mL (100 µL/well) can bind Biotinylated Human PVRIG, Fc,Avitag with a linear range of 0.005-0.313 µg/mL. Immobilized Human Nectin-2, Mouse IgG2a Fc Tag at 10 µg/mL (100 µL/well) can bind Biotinylated Human PVRIG, Fc,Avitag with a linear range of 0.01-0.313 µg/mL.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag). The protein has a calculated MW of 41.9 kDa. The protein migrates as 50-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 1.0 EU per µg by the LAL method.
Purity	>95% as determined by SDS-PAGE. >90% as determined by SEC-MALS.
Storage	For long term, the product should be stored at lyophilized state at -20°C or lower. Please avoid repeated freeze-thaw cycles. This product is stable after at: -20°C to -70°C for 12 months in lyophilized state; -70°C for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
Conjugation	Biotin

GENE INFORMATION

Gene Name	PVRIG poliovirus receptor related immunoglobulin domain containing [Homo sapiens]
Official Symbol	PVRIG
Synonyms	PVRIG; poliovirus receptor related immunoglobulin domain containing; transmembrane protein PVRIG; C7orf15; MGC2463; poliovirus receptor-related immunoglobulin domain-containing protein; MGC104322; MGC138295; MGC138297;
Gene ID	79037

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



mRNA Refseq [NM_024070](#)

Protein Refseq [NP_076975](#)

UniProt ID [Q6DKI7](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA