

Recombinant Human PVRL1 cell lysate

Cat. No. PVRL1-2299HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human PVRL1 / HVEC / CD111 / Nectin-1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

PVRL1 poliovirus receptor-related 1 (herpesvirus entry mediator C) [Homo sapiens]

Official Symbol

PVRL1

Synonyms

PVRL1; poliovirus receptor-related 1 (herpesvirus entry mediator C); ED4, HVEC; poliovirus receptor-related protein 1; CD111; CLPED1; HIgR; nectin; OFC7; PRR; PRR1; PVRR1; SK 12; nectin 1; poliovirus receptor-like 1; herpesvirus Ig-like receptor; herpes virus entry mediator C; ectodermal dysplasia 4 (Margarita Island type); ED4; HVEC; PVRR; SK-12; nectin-1; MGC16207; MGC142031;

Gene ID

5818

mRNA Refseq

NM_002855

Protein Refseq

NP_002846

MIM

600644

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q15223
Chromosome Location	11q23-q24
Pathway	Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Adherens junctions interactions, organism-specific biosystem; Cell adhesion molecules (CAMs), organism-specific biosystem; Cell adhesion molecules (CAMs), conserved biosystem; Cell junction organization, organism-specific biosystem; Cell-Cell communication, organism-specific biosystem;
Function	carbohydrate binding; cell adhesion molecule binding; coreceptor activity; protein binding; protein heterodimerization activity; protein homodimerization activity; receptor activity; virion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA